

MethoCult™ M3134



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Base methylcellulose medium for mouse cells

Catalog #03134

40 mL

Product Description

Base Methylcellulose Medium for Colony-Forming Unit (CFU) Assays for Mouse Cells

MethoCult™ M3134 is a base medium that contains 2.6% methylcellulose in Iscove's MDM. It does not contain serum, albumin, cytokines, or other medium supplements.

MethoCult™ M3134 is intended for making up a complete methylcellulose-based medium for CFU assays of mouse cells with the addition of desired components by the user. It is supplied in a volume of 40 mL per bottle. A 1% concentration of methylcellulose is obtained when brought to a volume of 100 mL.

Properties

Storage: Store at -20°C.

Shelf Life: Stable until expiry date (EXP) on label.

Contains:

- 2.6% Methylcellulose
- Iscove's MDM

Directions for Use

NOTE: If product is received partially thawed, place immediately at -20°C or thaw completely and aliquot as described below. Do not use MethoCult™ past the expiry date as indicated on the label.

PREPARATION OF COMPLETE METHOCULT™ M3134 MEDIUM

MethoCult™ M3134 base medium does not contain cytokines or other medium supplements; they can be added directly to the bottle or to each tube after aliquoting. Refer to Table 1 for volumes required to prepare complete MethoCult™ M3134 medium per bottle or per tube. The 2:3 (v:v) ratio of MethoCult™ to other components in the liquid medium (e.g. cytokines) gives the correct viscosity for optimal CFU growth and morphology.

Use sterile technique to prepare complete MethoCult™ M3134 medium (MethoCult™ M3134 base medium + desired components).

NOTE: Do not use pipettes to dispense methylcellulose as the volume dispensed will not be accurate. Syringes and large-bore, blunt-end needles should be used for accurate dispensing of viscous methylcellulose medium and to prevent needle-stick injuries.

A. TO PREPARE 100 mL BOTTLE

1. Thaw 40 mL bottle of MethoCult™ M3134 at room temperature (15 - 25°C) or overnight at 2 - 8°C.

NOTE: Do not thaw MethoCult™ at 37°C.

2. Prepare desired components, including fetal bovine serum (FBS), bovine serum albumin (BSA), cytokines, and Iscove's Modified Dulbecco's Medium (IMDM; Catalog #36150), as required, in 60 mL and add to MethoCult™ (total volume of 100 mL).
3. Shake vigorously for 1 - 2 minutes and then let stand for at least 5 minutes to allow bubbles to rise to the top before aliquoting.
4. Using a 3 or 6 mL luer lock syringe attached to a 16 Gauge Blunt-End Needle (Catalog #28110), aliquot 3 mL per tube for 1.1 mL duplicate cultures, or 4 mL per tube for 1.1 mL triplicate cultures. Complete MethoCult™ medium is now ready for use.

B. TO PREPARE INDIVIDUAL TUBES

1. Thaw 40 mL bottle of MethoCult™ M3134 at room temperature (15 - 25°C) or overnight at 2 - 8°C.

NOTE: Do not thaw MethoCult™ at 37°C.

2. Shake vigorously for 1 - 2 minutes and then let stand for at least 5 minutes to allow bubbles to rise to the top before aliquoting.

3. Using a 3 or 6 mL luer lock syringe attached to a 16 Gauge Blunt-End Needle (Catalog #28110), aliquot MethoCult™ M3134 base medium into tubes (see Table 1 for required volumes).
NOTE: Aliquots of MethoCult™ medium may be stored at -20°C before adding components, until expiry date as indicated on label. After thawing aliquots, add desired components and mix well.
4. Add desired growth factors, supplements, and Iscove's Modified Dulbecco's Medium (IMDM; Catalog #36150) to tubes of MethoCult™ M3134 (see Table 1 for required volumes).
5. Vortex tubes to mix well. Complete MethoCult™ medium is now ready for use.
6. Aliquot any remaining MethoCult™ M3134 base medium for duplicate or triplicate cultures (see Table 1 for required volumes), store at -20°C, then add desired components after thawing. Mix well before use.

Table 1. Volumes Required for Preparation of Complete MethoCult™ M3134 Medium

COMPONENT	PER BOTTLE	PER TUBE (duplicate 1.1 mL cultures)	PER TUBE (triplicate 1.1 mL cultures)
MethoCult™ M3134	40 mL	1.2 mL	1.6 mL
IMDM with cytokines*	60 mL	1.8 mL	2.4 mL
TOTAL VOLUME	100 mL	3 mL	4 mL

*For a complete list of available cytokines, refer to www.stemcell.com.

For recommended cell plating concentrations, setup of mouse CFU assays, and counting and classification of mouse colonies, refer to the Technical Manual: Mouse Colony-Forming Cell Assays Using MethoCult™ (Document #10000005597), available at www.stemcell.com, or contact us to request a copy.

References

Miller CL & Lai B. (2005) Human and mouse hematopoietic colony-forming cell assays. In: Helgason CD & Miller CL (Eds.). Basic Cell Culture Protocols (pp. 71–89). Totowa, New Jersey: Humana Press Inc.

Related Products

For related products, including specialized culture and storage media, supplements, antibodies, cytokines, and small molecules, visit www.stemcell.com/HSPCworkflow, or contact us at techsupport@stemcell.com.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2026 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and MethoCult are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.