

ClonaCell™-HY Medium B

Hybridoma fusion medium (serum-free)

Catalog #03802

500 mL



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

ClonaCell™-HY Medium B is a serum-free liquid medium used to wash lymphocytes and parental myeloma cells to remove serum prior to fusion and during the fusion process. This medium has been verified for use in mouse and rat hybridoma development and reportedly is compatible for production of hybridomas using lymphocytes from a variety of host animals including human, mouse, rat, and hamster.

Properties

Storage:	Store at -20°C.
Shelf Life:	Stable until expiry date (EXP) on label.
Contains:	<ul style="list-style-type: none">• DMEM• Gentamicin• 2-Mercaptoethanol• Phenol red• L-Glutamine and other supplements• Other ingredients

Directions for Use

1. Thaw ClonaCell™-HY Medium B at room temperature (15 - 25°C) or overnight at 2 - 8°C. Mix well.
NOTE: Do not thaw ClonaCell™-HY Medium B in a 37°C water bath.
2. If ClonaCell™-HY Medium B is not used immediately, store at 2 - 8°C for up to 4 months. Alternatively, aliquot and store at -20°C until expiry date as indicated on the label.

For further information, refer to the Technical Manual: ClonaCell™-HY: A Complete Workflow for Hybridoma Generation (Document #10000005590), available at www.stemcell.com, or contact us to request a copy.

References

- Chang Q et al. (2002) Structure-function relationships for human antibodies to pneumococcal capsular polysaccharide from transgenic mice with human immunoglobulin loci. *Infect Immun* 70(9): 4977–86.
- Chen ZC et al. (2000) Genes coding evolutionary novel anti-carbohydrate antibodies: Studies on anti-Gal production in alpha 1,3galactosyltransferase knock out mice. *Mol Immunol* 37(8): 455–66.
- Fang L et al. (2008) Essential role of TNF receptor superfamily 25 (TNFRSF25) in the development of allergic lung inflammation. *J Exp Med* 205(5): 1037–48.
- Flyak AI et al. (2015) Mechanism of human antibody-mediated neutralization of Marburg virus. *Cell* 160(5): 893–903.
- Kaabinejadian S et al. (2016) Immunodominant West Nile virus T cell epitopes are fewer in number and fashionably late. *J Immunol* 196(10): 4263–73.
- Kuroki M et al. Preparation of human IgG and IgM monoclonal antibodies for MK-1/Ep-CAM by using human immunoglobulin genetransferred mouse and gene cloning of their variable regions. *Anticancer Res* 25(6A): 3733–9.
- Loveless BC et al. (2011) Structural characterization and epitope mapping of the glutamic acid/alanine-rich protein from *Trypanosoma congolense*: Defining assembly on the parasite cell surface. *J Biol Chem* 286(23): 20658–65.
- Okai S et al. (2016) High-affinity monoclonal IgA regulates gut microbiota and prevents colitis in mice. *Nat Microbiol* 1(9): 16103.
- Retamal M et al. (2014) Epitope mapping of the 2009 pandemic and the A/Brisbane/59/2007 seasonal (H1N1) influenza virus haemagglutinins using mAbs and escape mutants. *J Gen Virol* 95(11): 2377–89.
- Smith SA et al. (2012) Persistence of circulating memory B cell clones with potential for Dengue virus disease enhancement for decades following infection. *J Virol* 86(5): 2665–75.

- Spanier JA et al. (2016) Efficient generation of monoclonal antibodies against peptide in the context of MHCII using magnetic enrichment. *Nat Commun* 7: 11804.
- Wang X et al. (2013) Generation and characterization of a unique reagent that recognizes a panel of recombinant human monoclonal antibody therapeutics in the presence of endogenous human IgG. *MAbs* 5(4): 540–54.
- Wittman VP et al. (2006) Antibody targeting to a class I MHC-peptide epitope promotes tumor cell death. *J Immunol* 177(6): 4187–95.
- Yew CW & Tan YJ. (2016) Generation of mouse monoclonal antibodies specific to chikungunya virus using ClonaCell-HY Hybridoma Cloning Kit. *Methods Mol Biol* 1426: 225–33.
- Yuki N et al. (2004) Carbohydrate mimicry between human ganglioside GM1 and *Campylobacter jejuni* lipooligosaccharide causes Guillain-Barre syndrome. *Proc Natl Acad Sci USA* 101(31): 11404–9.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2026 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and ClonaCell are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.