

# Anti-Human CD32 Antibody, Clone IV.3, FITC

Mouse monoclonal IgG2b antibody against human CD32, FITC-conjugated

Catalog #60012Fl.1 25 Tests 20 µL/test

Catalog #100-1574 100 Tests 20 µL/test

## Product Description

This monoclonal antibody reacts with human CD32 (FcγRII), an ~40 kDa type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, immunomodulation and platelet aggregation. CD32 is encoded by three genes (A, B, C) and at least 6 isoforms are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 and IIc. All isoforms are expressed by monocytes/macrophages, placental trophoblasts and endothelial cells. In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes and, weakly, by B cells. Isoform IIc is expressed by NK cells and neutrophils. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexes. These interactions can result in non-specific labeling in antibody-based detection and cell separation experiments and the IV.3 antibody may be employed as a blocking antibody to reduce non-specific binding. The IV.3 antibody binds most strongly to the IIa isoforms of CD32, with the epitope mapped to amino acids 132 - 137 [FSHLDP] in domain 2, within the ligand binding site. Binding of the IV.3 antibody can be blocked by clone FLI8.26 in flow cytometry analyses, suggesting that these clones may share a common or overlapping epitope.

Target Antigen:	CD32
Alternative Names:	FCR II, FcγRII
Gene ID:	2212
Species Reactivity:	Human
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	IV.3
Isotype:	IgG2b, kappa
Immunogen:	K-562 human erythromyeloblastoid leukemia cell line
Conjugate:	FITC (Fluorescein isothiocyanate)

## Applications

**Verified Applications:** CellSep, FC

**Reported Applications:** FC

**Special Applications:** This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human T Cell Enrichment Kit (Catalog #19051) and EasySep™ Human CD4+ T Cell Enrichment Kit (Catalog #19052).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; FCXM: Flow cytometric crossmatch assay; FISH: Fluorescence in situ hybridization; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IHC-F: Immunohistochemistry (frozen-tissue); IHC-P: Immunohistochemistry (paraffin-embedded); IP: Immunoprecipitation; NMR: Nuclear magnetic resonance spectroscopy; RIA: Radioimmunoassay; WB: Western blotting

## Properties


**Product Formulation:** Phosphate-buffered saline containing less than 0.1% (w/v) sodium azide and less than 0.1% (w/v) bovine serum albumin

**Purification:** The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions.

**Stability and Storage:** Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, contact [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

**Directions for Use:** For flow cytometry, the suggested use of this antibody is 20 µL per  $1 \times 10^6$  cells in 100 µL or per 100 µL of whole blood. It is recommended that the antibody be titrated for optimal performance for each application.

## Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD32 Antibody, Clone IV.3, FITC (filled histogram) or Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11, FITC (Catalog #60072FI; solid line histogram).

## Related Products

For a complete list of antibodies, including other conjugates, sizes, and clones, as well as related products available from STEMCELL Technologies, visit [www.stemcell.com/antibodies](http://www.stemcell.com/antibodies), or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

- Boruchov AM et al. (2005) Activating and inhibitory IgG Fc receptors on human DCs mediate opposing functions. *J Clin Invest* 115(10): 2914–23.
- Ierino FL et al. (1993) Mapping epitopes of human Fc gamma RII (CDw32) with monoclonal antibodies and recombinant receptors. *J Immunol* 150(5): 1794–803.
- Joulia R et al. (2015) Mast cells form antibody-dependent degranulatory synapse for dedicated secretion and defence. *Nat Commun* 6: 6174.
- Looney RJ et al. (1986) Human monocytes and U937 cells bear two distinct Fc receptors for IgG. *J Immunol* 136(5): 1641–7.
- Micklem KJ et al. (1990) Different isoforms of human FcRII distinguished by CDw32 antibodies. *J Immunol* 144(6): 2295–303.
- Ramsland PA et al. (2011) Structural basis for FcγRIIa recognition of human IgG and formation of inflammatory signaling complexes. *J Immunol* 187(6): 3208–17.
- Sardjono CT et al. (2008) Epitope mapping of Fc gamma RIIa monoclonal antibodies. *Indonesian J Biotechnol* 13(1): 1030–7.
- Schlossman SF et al. (Eds.). (1995) Binding heterogeneity within the CD32 panel of mAB. In: *Leucocyte Typing V* (pp. 832-35). New York: Oxford University Press.
- Tomiyama Y et al. (1992) Response of human platelets to activating monoclonal antibodies: Importance of Fc gamma RII (CD32) phenotype and level of expression. *Blood* 80(9): 2261–8.
- van Sorge NM et al. (2003) FcγRII polymorphisms: Implications for function, disease susceptibility and immunotherapy. *Tissue Antigens* 61(3): 189–202.
- Vely F et al. (1997) A new set of monoclonal antibodies against human Fc gamma RII (CD32) and Fc gamma RIII (CD16): characterization and use in various assays. *Hybridoma* 16(6): 519–28.
- Xiao Z et al. (2008) Immune complexes formed following the binding of anti-platelet factor 4 (CXCL4) antibodies to CXCL4 stimulate human neutrophil activation and cell adhesion. *Blood* 112(4): 1091–100.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.