

# Ascorbic Acid

Antioxidant; Reducing agent

Catalog #72132 500 mg

Catalog #100-1040 100 g

## Product Description

Ascorbic Acid is a naturally occurring lactone that is produced by plants and many animals, but not humans or other primates. It acts as an electron donor (i.e. reducing agent) and shows antioxidant activity particularly against reactive oxygen species. Ascorbic Acid is a cofactor for monooxygenase and dioxygenase as well as other enzymes (Arrigoni & De Tullio; Du et al.).

**Molecular Name:** Ascorbic Acid

**Alternative Names:** (+)-Ascorbic Acid; L-Ascorbic Acid; NSC 218455; NSC 33832; Vitamin C

**CAS Number:** 50-81-7


**Chemical Formula:** C<sub>6</sub>H<sub>8</sub>O<sub>6</sub>

**Molecular Weight:** 176.1 g/mol

**Purity:** ≥ 95%

**Chemical Name:** (R)-5-((S)-1,2-dihydroxyethyl)-3,4-dihydroxyfuran-2(5H)-one

**Structure:**



## Properties

|                               |   |
|-------------------------------|---|
| <b>Product Format:</b>        | A crystalline solid   |
| <b>Stability and Storage:</b> | Product stable at -20°C as supplied. As a precaution, STEMCELL recommends storing all small molecules away from direct light. Stable as supplied for 12 months from date of receipt.  |
| <b>Preparation:</b>           | <ul style="list-style-type: none"><li>• PBS (pH 7.2) ≤ 85 mM</li><li>• Water ≤ 85 mM</li></ul> <p>For example, to prepare a 10 mM stock solution in PBS, resuspend 100 mg in 56.8 mL of PBS (pH 7.2). Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported; however, as a general guide we recommend storage in PBS at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.</p> <p>For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use.</p> |

## Published Applications

### MAINTENANCE AND SELF-RENEWAL

- Supports proliferation of mesenchymal stem cells (Choi et al.).

### REPROGRAMMING

- Increases the efficiency of reprogramming mouse and human fibroblasts to induced pluripotent stem (iPS) cells (Esteban et al.) partly through JHDM1 histone demethylase activity (Wang et al.).
- Prevents aberrant DNA methylation of the Dlk1-Dio3 locus during reprogramming of mouse somatic cells to iPS cells (Stadtfield et al.)

### DIFFERENTIATION

- Promotes differentiation of osteoblasts from human and mouse mesenchymal cells (Pittenger et al.; Tropel et al.).
- Promotes differentiation of osteoblasts from mouse embryonic stem (ES) cells (zur Nieden et al.).
- Enhances differentiation of cardiomyocytes from mouse ES cells (Takahashi et al.).

## References

- Arrigoni O & De Tullio MC. (2002) Ascorbic acid: much more than just an antioxidant. *Biochim Biophys Acta* 1569(1-3): 1–9.
- Choi K-M et al. (2008) Effect of ascorbic acid on bone marrow-derived mesenchymal stem cell proliferation and differentiation. *J Biosci Bioeng* 105(6): 586–94.
- Du J et al. (2012) Ascorbic acid: chemistry, biology and the treatment of cancer. *Biochim Biophys Acta* 1826(2): 443–57.
- Esteban MA et al. (2010) Vitamin C enhances the generation of mouse and human induced pluripotent stem cells. *Cell Stem Cell* 6(1): 71–9.
- Pittenger MF et al. (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284(5411): 143–7.
- Stadtfield M et al. (2012) Ascorbic acid prevents loss of Dlk1-Dio3 imprinting and facilitates generation of all-iPS cell mice from terminally differentiated B cells. *Nat Genet* 44(4): 398–405, S1–2.
- Takahashi T et al. (2003) Ascorbic acid enhances differentiation of embryonic stem cells into cardiac myocytes. *Circulation* 107(14): 1912–6.
- Tropel P et al. (2004) Isolation and characterisation of mesenchymal stem cells from adult mouse bone marrow. *Exp Cell Res* 295(2): 395–406.
- Wang T et al. (2011) The histone demethylases Jhdm1a/1b enhance somatic cell reprogramming in a vitamin-C-dependent manner. *Cell Stem Cell* 9(6): 575–87.
- Zur Nieden NI et al. (2003) In vitro differentiation of embryonic stem cells into mineralized osteoblasts. *Differentiation* 71(1): 18–27.

## Related Products

For a complete list of small molecules available from STEMCELL Technologies, visit [www.stemcell.com/smallmolecules](http://www.stemcell.com/smallmolecules) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## Ascorbic Acid

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.