

Small Molecules

Reversine

Adenosine receptor, non-muscle myosin II (NM II), MEK1, and aurora kinase inhibitor

Catalog # 72612
72614

1 mg
5 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Reversine is a 2,6-disubstituted purine derivative that was identified using a screening system based on the loss of terminal differentiation markers in C2C12 myoblast cells (Chen et al. 2004). Reversine has been shown to inhibit the human A3 adenosine receptor, nonmuscle myosin II heavy chain, mitogen-activated extracellular signal-regulatedkinase-1 (MEK1), and Aurora B kinase (Chen et al. 2007; Perreira et al.; D'Alise et al.).

Molecular Name:	Reversine
Alternative Names:	Not applicable
CAS Number:	656820-32-5
Chemical Formula:	C ₂₁ H ₂₇ N ₇ O
Molecular Weight:	393.5 g/mol
Purity:	> 95%
Chemical Name:	N6-cyclohexyl-N2-[4-(4-morpholinyl)phenyl]-1H-purine-2,6-diamine
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. Stable as supplied for 12 months from date of receipt.
Solubility:	· DMSO ≤ 25 mM For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 254 µL of DMSO. Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported; however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application. Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

REPROGRAMMING

- Induces dedifferentiation of lineage-committed murine myoblasts to multipotent progenitors with osteogenic and adipogenic potential (Chen et al., 2007; Chen et al., 2004).
- Induces dedifferentiation of primary mouse and human dermal fibroblasts to myogenic-competent cells (Anastasia et al.).
- Induces dedifferentiation of annulus fibrosus cells to multipotent mesenchymal progenitors that have the potential to develop along chondrogenic, osteogenic, or adipogenic lineages (Saraiya et al.).

CANCER RESEARCH

- Inhibits cell proliferation and induces polyploidy in the PC-3, HeLa, CWR22Rv1, and DU-145 cancer cell lines (Hsieh et al.).
- Induces differentiation in the human embryonal carcinoma cell line NT2, and in the human promyelocytic leukemia cell line HL60 (D'Alise et al.).
- Blocks proliferation, causes failure of cytokinesis, and induces polyploidy in multiple cancer cell lines (D'Alise et al.).

References

- Anastasia L et al. (2006) Reversine-treated fibroblasts acquire myogenic competence in vitro and in regenerating skeletal muscle. *Cell Death Differ* 13(12): 2042–51.
- Chen S et al. (2004) Dedifferentiation of lineage-committed cells by a small molecule. *J Am Chem Soc* 126(2): 410–1.
- Chen S et al. (2007) Reversine increases the plasticity of lineage-committed mammalian cells. *Proc Natl Acad Sci USA* 104(25): 10482–7.
- D'Alise AM et al. (2008) Reversine, a novel Aurora kinases inhibitor, inhibits colony formation of human acute myeloid leukemia cells. *Mol Cancer Ther* 7(5): 1140–9.
- Hsieh T-C et al. (2007) The 2,6-disubstituted purine reversine induces growth arrest and polyploidy in human cancer cells. *Int J Oncol* 31(6): 1293–300.
- Perreira M et al. (2005) “Reversine” and its 2-substituted adenine derivatives as potent and selective A3 adenosine receptor antagonists. *J Med Chem* 48(15): 4910–8.
- Saraiya M et al. (2010) Reversine enhances generation of progenitor-like cells by dedifferentiation of annulus fibrosus cells. *Tissue Eng Part A* 16(4): 1443–55.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2022 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.