

Small Molecules

Lapatinib

Tyrosine kinase inhibitor; Inhibits EGFR and HER2

Catalog # 73242
73244

10 mg
100 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Lapatinib is a selective dual inhibitor of epidermal growth factor receptor (EGFR, ErbB1) and human epidermal growth factor receptor 2 (HER2, ErbB2), each with an IC_{50} of about 10 nM (Rusnak et al.). It binds reversibly, but with a very slow off-rate, to the ATP binding pocket in an inactive-like conformation, thus preventing autophosphorylation. Binding is very potent, with K_i values of 3 and 13 nM to EGFR and HER2, respectively (Wood et al.).

Molecular Name: Lapatinib
Alternative Names: GSK 572016; GW 572016; GW2016; Tyverb
CAS Number: 231277-92-2
Chemical Formula: $C_{29}H_{26}ClF_4O_4S$
Molecular Weight: 581.1 g/mol
Purity: $\geq 98\%$
Chemical Name: N-[3-chloro-4-[(3-fluorophenyl)methoxy]phenyl]-6-[5-[[[2-(methylsulfonyl)ethyl]amino]methyl]-2-furanyl]-4-quinazolinamine
Structure:



Properties

Physical Appearance: A crystalline solid
Storage: Product stable at -20°C as supplied. Protect from prolonged exposure to light.
Stable as supplied for 12 months from date of receipt.
Solubility: · DMSO ≤ 30 mM
For example, to prepare a 10 mM stock solution in DMSO, resuspend 10 mg in 1.72 mL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

CANCER RESEARCH

- Induces apoptosis in cells overexpressing EGFR and/or HER2, and inhibits tumor growth in a mouse xenograft model (Xia et al.).

References

Rusnak DW et al. (2001) The effects of the novel, reversible epidermal growth factor receptor/ErbB-2 tyrosine kinase inhibitor, GW2016, on the growth of human normal and tumor-derived cell lines in vitro and in vivo. *Mol Cancer Ther* 1(2): 85–94.

Wood ER et al. (2004) A unique structure for epidermal growth factor receptor bound to GW572016 (Lapatinib): relationships among protein conformation, inhibitor off-rate, and receptor activity in tumor cells. *Cancer Res* 64(18): 6652–9.

Xia W et al. (2002) Anti-tumor activity of GW572016: a dual tyrosine kinase inhibitor blocks EGF activation of EGFR/erbB2 and downstream Erk1/2 and AKT pathways. *Oncogene* 21(41): 6255–63.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.