

# Human Recombinant bFGF

Basic fibroblast growth factor

Catalog #78003.1	10 µg
Catalog #78003	50 µg
Catalog #78003.2	1000 µg

## Product Description

Basic fibroblast growth factor (bFGF) is a prototypic member of the fibroblast growth factor family. Cytokines in the FGF family possess broad mitogenic and cell survival activities (Folkman & Klagsbrun; Kimelman & Kirschner) and are involved in a variety of biological processes including cell proliferation, differentiation, survival, and apoptosis (Folkman & Klagsbrun; Klagsbrun; Rifkin & Moscatelli). bFGF has the  $\beta$ -trefoil structure (Ponting & Russell), binds to the four FGF receptor (FGFR) family members, and activates JAK/STAT, PI3K, ERK1/2, and other receptor tyrosine kinase (RTK) signaling pathways. It supports the maintenance of undifferentiated human embryonic stem cells (Kang et al.; Xu et al.), stimulates human embryonic stem cells to form neural rosettes (Zhang et al.), improves proliferation of human mesenchymal stem cells, and enhances chondrogenic differentiation (Solchaga et al.).

Product Information

Alternative Names:	Basic fibroblast growth factor, FGF-β, FGF2, Fibroblast growth factor-basic, HBGF-2
Accession Number:	P09038
Amino Acid Sequence:	AAGSITTLPA LPEDGGSGAF PPGHFKDPKR LYCKNGGFFL RIHPDGRVDG VREKSDPHIK LQLQAEERGVS VSIKGVCANR YLAMKEDGRL LASKCVTDEC FFFERLESNN YNTYRSRKYT SWYVALKRTG QYKLGSKTGP GQKAILFLPM SAKS
Predicted Molecular Mass:	17.1 kDa
Species:	Human
Product Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 4 \times 10^6$ units/mg ( $EC_{50} \leq 0.25$ ng/mL) as determined by a cell proliferation assay using BALB/c 3T3 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is $\leq 0.2$ EU/μg protein.

Preparation and Storage

Stability and Storage:	Store at -80°C. Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. Store at 2 - 8°C for up to 2 weeks or at -20 to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.

Data



**Figure 1. Biological Activity of Human Recombinant bFGF**

(A) Biological activity of Human Recombinant bFGF was tested by its ability to promote the proliferation of BALB/c 3T3 cells. The EC<sub>50</sub> is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC<sub>50</sub> in the above example is 0.1 ng/mL. (B) 2 μg of Human Recombinant bFGF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant bFGF has a predicted molecular mass of 17.1 kDa.

## Related Products

For a complete list of cytokines or peptide pools, as well as related products available from STEMCELL Technologies, visit [www.stemcell.com/cytokines](http://www.stemcell.com/cytokines) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

- Folkman J & Klagsbrun M. (1987) Angiogenic factors. *Science* 235(4787): 442–7.
- Kang HB et al. (2005) Basic fibroblast growth factor activates ERK and induces c-fos in human embryonic stem cell line MizhES1. *Stem Cells Dev* 14(4): 395–401.
- Kimelman D & Kirschner M. (1987) Synergistic induction of mesoderm by FGF and TGF-beta and the identification of an mRNA coding for FGF in the early *Xenopus* embryo. *Cell* 51(5): 869–77.
- Klagsbrun M. (1989) The fibroblast growth factor family: structural and biological properties. *Prog Growth Factor Res* 1(4): 207–35.
- Ponting CP & Russell RB. (2000) Identification of distant homologues of fibroblast growth factors suggests a common ancestor for all beta-trefoil proteins. *J Mol Biol* 302(5): 1041–7.
- Rifkin DB & Moscatelli D. (1989) Recent developments in the cell biology of basic fibroblast growth factor. *J Cell Biol* 109(1): 1–6.
- Solchaga LA et al. (2005) FGF-2 enhances the mitotic and chondrogenic potentials of human adult bone marrow-derived mesenchymal stem cells. *J Cell Physiol* 203(2): 398–409.
- Xu RH et al. (2005) Basic FGF and suppression of BMP signaling sustain undifferentiated proliferation of human ES cells. *Nat Methods* 2(3): 185–90.
- Zhang SC et al. (2001) In vitro differentiation of transplantable neural precursors from human embryonic stem cells. *Nat Biotechnol* 19(12): 1129–33.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.