

Cytokines

Human/Mouse Recombinant BMP-2, ACF



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	78135	10 µg
	78135.1	100 µg
	78135.2	1000 µg

Product Description

Bone morphogenetic protein 2 (BMP-2) is a member of the transforming growth factor beta (TGF- β) superfamily. BMP-2 is a disulfide-linked homodimer, acts as a ligand for complexes of type I and II BMP receptors, and primarily activates SMAD1/5/8 signaling (Nohe et al.). BMP-2 is a potent differentiation factor and directs human pluripotent stem cells (hPSCs) towards various cell types including extra-embryonic endoderm, mesenchymal cells, and chondrocytes (Pera et al.). Although BMP-2 expression is low in healthy cartilage, its expression is upregulated at the site of cartilage damage (Blaney Davidson et al.). BMP-2 induces bone and cartilage formation in vitro and is able to induce chondrogenesis in human mesenchymal stem cells (Schmitt et al.). This product is animal component-free.

Product Information

Alternative Names:	BDA2, BMP-2A, BMP2, Bone morphogenetic protein 2A
Accession Number:	P12643
Amino Acid Sequence:	MQAKHKQRKR LKSSCKRHPV YVDFSDVGWN DWIVAPPGYH AFYCHGECPF PLADHLNSTN HAIQQLVNS VNSKIPKACC VPTELSAISM LYLDENEKVV LKNYQDMVVE GCGCR
Predicted Molecular Mass:	13 kDa monomer; 26.1 kDa dimer
Species:	Human, Mouse
Formulation:	Lyophilized from a sterile-filtered aqueous solution containing 0.1% trifluoroacetic acid.
Source:	<i>E. coli</i>

Specifications


Activity:	The specific activity is $\geq 4.0 \times 10^3$ units/mg (EC50 ≤ 250 ng/mL) as determined by alkaline phosphatase activity induced in ATDC-5 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/ μ g protein.

Preparation and Storage

Storage:	Store at -20°C to -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide we recommend to not store at 2 - 8°C for more than 1 month or at -20°C to -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human/Mouse Recombinant BMP-2, ACF was tested by its ability to induce alkaline phosphatase activity in ATDC-5 cells. The EC50 is defined as the effective concentration of the growth factor at which alkaline phosphatase production is at 50% of maximum. The EC50 in the above example is 74 ng/mL.

(B) 1 µg of Human/Mouse Recombinant BMP-2, ACF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human/Mouse Recombinant BMP-2, ACF has a predicted molecular mass of 13 kDa (monomer) or 26.1 kDa (dimer).

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Blaney Davidson EN et al. (2007) Elevated extracellular matrix production and degradation upon bone morphogenetic protein-2 (BMP-2) stimulation point toward a role for BMP-2 in cartilage repair and remodeling. *Arthritis Res Ther* 9(5): R102.

Nohe A et al. (2001) The mode of bone morphogenetic protein (BMP) receptor oligomerization determines different BMP-2 signaling pathways. *J Bio Chem* 277: 5330–8.

Pera MF et al. (2004) Regulation of human embryonic stem cell differentiation by BMP-2 and its antagonist noggin. *J Cell Sci* 117(Pt 7): 1269–80.

Schmitt B et al. (2003) BMP2 initiates chondrogenic lineage development of adult human mesenchymal stem cells in high-density culture. *Differentiation* 71(9-10): 567–77.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2021 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.