

Human Recombinant R-Spondin-1 (CHO-expressed)

R-Spondin-1

Catalog #78213	25 µg
Catalog #78213.1	100 µg
Catalog #78213.2	500 µg
Catalog #78213.3	1000 µg

Product Description

R-Spondin-1 (RSPO1) is the prototype member of the R-Spondin (RSPO) protein subfamily of a superfamily of thrombospondin type 1 repeat (TSR-1)-containing proteins (Chen et al.; Kamala et al.; Kazanskaya et al.; Kim et al.). Although unable to initialize signaling, RSPO family members are potent enhancers of WNT signaling (Cruciat & Niehrs; de Lau et al.; Kamala et al.; Kazanskaya et al.). They are characterized by a TSR-1 domain, a carboxy-terminal region with positively charged amino acids, and two N-terminal furin-like cysteine-rich repeats (Glinka et al.; Kazanskaya et al.). RSPO1 activates β -catenin signaling via the WNT signaling cascade and by indirectly increasing low-density lipoprotein receptor-related protein 6 (LRP6) on the cell surface. It does this by binding leucine-rich repeat-containing G-protein-coupled receptor 5 (LGR5), and competing with WNT antagonist DKK-1 for binding to the WNT coreceptors Kremen and LRP6, which reduces DKK-1-mediated internalization of LRP6 (Binnerts et al.). RSPO1 is involved in a wide range of pleiotropic roles during embryogenesis, it is required for the specification of hematopoietic stem cells, and it has been shown to be important in the growth, survival, and migration of ovarian cancer cells (Cruciat & Niehrs; de Lau et al.; Genthe & Clements; Liu et al.).

Product Information

Alternative Names:	hRspo1, Roof plate-specific Spondin-1
Accession Number:	Q2MKA7
Amino Acid Sequence:	SRGIKGRQR RISAEGSQAC AKGCELCSEV NGCLKCSPKL FILLERNDIR QVGVCLPSCP PGYFDARNPD MNKCIKCKIE HCEACFSHNF CTKCKEGLYL HKGRCYPACP EGSSAANGTM ECSSPAQCEM SEWSPWGPCS KKQQLCGFRR GSEERTRRVL HAPVGDHAAC SDKETRRCCT VRRVPCPEGQ KRRKGGQGRR ENANRNLARK ESKEAGAGSR RRKGGQQQQQ QGTVGPLTSA GPA
Predicted Molecular Mass:	26.8 kDa
Species:	Human
Product Formulation:	Lyophilized from a sterile-filtered solution containing phosphate-buffered saline.
Source:	CHO
Purity:	≥ 95%


Specifications

Activity:	The specific activity is $\geq 2.0 \times 10^4$ units/mg ($EC_{50} \leq 50$ ng/mL), as determined by luciferase activity induced in HEK-293 cells in the presence of mouse Wnt-3a.
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/ μ g protein.

Preparation and Storage

Stability and Storage:	Store at -20 to -80°C. Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -20 to -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant R-Spondin-1 was tested by its ability to induce luciferase activity in a WNT reporter HEK-293 cell line in the presence of mouse Wnt-3a. Luciferase activity was measured using a luminometric assay method. The EC50 is defined as the effective concentration of the growth factor at which luciferase activity is at 50% of maximum. The EC50 in the above example is 36.4 ng/mL. (B) Human Recombinant R-Spondin-1 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant R-Spondin-1 has a predicted molecular mass of 26.8 kDa.

Related Products

For a complete list of cytokines or peptide pools, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Binnerts ME et al. (2007) R-Spondin1 regulates Wnt signaling by inhibiting internalization of LRP6. *Proc Natl Acad Sci USA* 104(37): 14700–5.
- Chen JZ et al. (2002) Cloning and identification of a cDNA that encodes a novel human protein with thrombospondin type I repeat domain, hPWTSR. *Mol Biol Rep* 29(3): 287–92.
- Cruciat C-M & Niehrs C. (2013) Secreted and transmembrane wnt inhibitors and activators. *Cold Spring Harb Perspect Biol* 5(3): a015081.
- de Lau WBM et al. (2012) The R-spondin protein family. *Genome Biol* 13(3): 242.
- Genthe JR & Clements WK. (2017) R-spondin 1 is required for specification of hematopoietic stem cells through Wnt16 and Vegfa signaling pathways. *Development* 144(4): 590–600.
- Glinka A et al. (2011) LGR4 and LGR5 are R-spondin receptors mediating Wnt/ β -catenin and Wnt/PCP signalling. *EMBO Rep* 12(10): 1055–61.
- Kazanskaya O et al. (2004) R-Spondin2 is a secreted activator of Wnt/ β -catenin signaling and is required for *Xenopus* myogenesis. *Dev Cell* 7(4): 525–34.
- Kim KA et al. (2006) R-Spondin proteins: a novel link to β -catenin activation. *Cell Cycle* 5(1): 23–6.
- Liu Q et al. (2019) The role of R-spondin 1 through activating Wnt/ β -catenin in the growth, survival and migration of ovarian cancer cells. *Gene* 689: 124–30.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.