EasySep™ Mouse CD11b Positive Selection Kit II

For processing 2 x 10⁹ cells from bone marrow

Catalog #18970 Catalog #18970RF RoboSep™

Positive Selection

Document #10000005377 | Version 02



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713
INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Description

Isolate highly purified CD11b+ cells from mouse bone marrow samples by positive selection.

- · Fast, easy-to-use, and column-free
- Up to 99% purity
- · Isolated cells are not fluorochrome-labeled

This kit targets CD11b+ cells for positive selection with antibodies recognizing the CD11b surface marker. Desired cells are labeled with antibodies and magnetic particles and separated without columns using an EasySep™ magnet. Unwanted cells are simply poured off, while desired cells remain in the tube. Isolated cells are immediately available for downstream applications such as flow cytometry, cell culture, and cell-based experiments.

NOTE: This is the Product Information Sheet (PIS) for isolating CD11b+ cells from mouse bone marrow samples. If isolating CD11b+ cells from mouse spleen, lung, or brain tissues, refer to the applicable PIS, available at www.stemcell.com, or contact us to request a copy.

Component Descriptions

| COMPONENT NAME | COMPONENT # | QUANTITY | STORAGE | SHELF LIFE | FORMAT |
|-----------------------------------------------------------------------|-------------|------------|-------------------------------------|------------------------------------------|----------------------------------------------------------------------------------------|
| EasySep [™] Mouse CD11b Positive Selection II Component A | 18970CA | 1 x 0.5 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A combination of monoclonal antibodies in PBS and 0.1% BSA. |
| EasySep™ Mouse CD11b Positive Selection II Component B | 18970CB | 1 x 0.5 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A combination of monoclonal antibodies in PBS and 0.1% BSA. |
| EasySep™ Dextran RapidSpheres™ 50100 | 50100 | 2 x 1 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A suspension of magnetic particles in water. |
| RoboSep™ Empty Vial | 27401 | 1 | Not applicable | Not applicable | Not applicable |
| Mouse FcR PolyBlock | 300-0902 | 1 x 1.2 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A combination of polyclonal antibodies and maltose in water with 5 µg/mL Triton X-100. |

BSA - bovine serum albumin; PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

Additional Reagent Stability Information

| REAGENT NAME | STORAGE | SHELF LIFE |
|---------------------------------------------------------|----------------------------------|-----------------------------------------------------------------------------------------|
| Selection Cocktail (combined Component A + Component B) | Store at 2 - 8°C. Do not freeze. | Stable for up to 4 weeks. Do not exceed the expiry date (EXP) of individual components. |

EasySep™ Mouse CD11b Positive Selection Kit II



Sample Preparation

BONE MARROW

Flush bone marrow cells from femur and tibia into recommended medium using a syringe equipped with a 23 gauge needle. Disperse clumps by gently passing the cell suspension through the syringe several times. Alternatively, crush bones using a mortar and pestle. Remove remaining clumps and debris by passing cell suspension through a 70 µm mesh nylon strainer (e.g. Catalog #27260). Centrifuge at 300 x g for 10 minutes and resuspend at 1 x 10^8 nucleated cells/mL in recommended medium.

SPLEEN, LUNG, OR BRAIN TISSUE

If processing spleen tissue, lung, or brain tissue, refer to the applicable PIS, available at www.stemcell.com, or contact us to request a copy.

Recommended Medium

EasySep™ Buffer (Catalog #20144), RoboSep™ Buffer (Catalog #20104), or PBS containing 2% FBS and 1 mM EDTA. Medium should be free of Ca++ and Mg++.



Directions for Use – Manual EasySep™ Protocols

See page 2 for Sample Preparation and Recommended Medium. Refer to Tables 1 and 2 for detailed instructions regarding the EasySep™ procedure for each magnet.

Table 1. EasySep™ Mouse CD11b Positive Selection Kit II Protocol for BONE MARROW

| | | EASYSEP™ MAGNETS | | | |
|------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|--|--|
| STEP | INSTRUCTIONS | EasySep™ (Catalog #18000) | "The Big Easy" (Catalog #18001) | | |
| 1 | Prepare sample at the indicated cell concentration within the volume range. | 1 x 10^8 cells/mL 0.25 - 2 mL | 1 x 10^8 cells/mL 1 - 4 mL | | |
| 2 | Add Mouse FcR PolyBlock to sample. | 50 μL/mL of sample | 50 μL/mL of sample | | |
| 3 | Add sample to required tube. | 5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Catalog #38007) | 14 mL (17 x 95 mm) polystyrene round-bottom tube (e.g. Catalog #38008) | | |
| 4 | Prepare Selection Cocktail in a tube. For each 1 mL of sample make 50 µL of cocktail (25 µL of Component A + 25 µL of Component B). | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | | |
| | Incubate. | RT for 5 minutes | RT for 5 minutes | | |
| 5 | Add Selection Cocktail to sample. NOTE: Do not vortex cocktail. | 50 μL/mL of sample | 50 μL/mL of sample | | |
| | Mix and incubate. | RT for 5 minutes | RT for 5 minutes | | |
| 6 | Vortex RapidSpheres™. NOTE: Particles should appear evenly dispersed. | 30 seconds | 30 seconds | | |
| 7 | Add RapidSpheres™ to sample. | 100 μL/mL of sample | 100 μL/mL of sample | | |
| 7 | Mix and incubate. | RT for 3 minutes | RT for 3 minutes | | |
| 8 | Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times. | Top up to 2.5 mL | Top up to 5 mL for samples < 3 mL Top up to 10 mL for samples ≥ 3 mL | | |
| | Place the tube (without lid) into the magnet and incubate. | RT for 3 minutes | RT for 5 minutes | | |
| 9 | Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells. | Discard supernatant | Discard supernatant | | |
| 10 | Repeat steps as indicated. | Steps 8 and 9, three more times (total of 4 x 3-minute separations) | Steps 8 and 9, three more times (total of 4 x 5-minute separations) | | |
| 11 | Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube. | Isolated cells are ready for use | Isolated cells are ready for use | | |
| | | | | | |

RT - room temperature (15 - 25°C)

^{*} Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.



Table 2. EasySep™ Mouse CD11b Positive Selection Kit II Protocol for BONE MARROW

| | | rotocol for BONE MARROW EASYSEP™ MAGNETS | | | |
|------|-------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|--|--|
| 211 | INSTRUCTIONS | EasyEights™ (Catalog #18103) | | | |
| STEP | | 5 mL tube | 14 mL tube | | |
| 1 | Prepare sample at the indicated cell concentration within the volume range. | 1 x 10^8 cells/mL 0.5 - 1.5 mL | 1 x 10^8 cells/mL 1 - 4 mL | | |
| 2 | Add Mouse FcR PolyBlock to sample. | 50 μL/mL of sample | 50 μL/mL of sample | | |
| 3 | Add sample to required tube. | 5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Catalog #38007) | 14 mL (17 x 95 mm) polystyrene round-bottom tube (e.g. Catalog #38008) | | |
| 4 | Prepare Selection Cocktail in a tube. For each 1 mL of sample make 50 μL of cocktail (25 μL of Component A + 25 μL of Component B). | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | | |
| | Incubate. | RT for 5 minutes | RT for 5 minutes | | |
| 5 | Add Selection Cocktail to sample. NOTE: Do not vortex cocktail. | 50 μL/mL of sample | 50 μL/mL of sample | | |
| | Mix and incubate. | RT for 5 minutes | RT for 5 minutes | | |
| 6 | Vortex RapidSpheres™. NOTE: Particles should appear evenly dispersed. | 30 seconds | 30 seconds | | |
| - | Add RapidSpheres™ to sample. | 100 μL/mL of sample | 100 μL/mL of sample | | |
| 7 | Mix and incubate. | RT for 3 minutes | RT for 3 minutes | | |
| 8 | Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times. | Top up to 2.5 mL | Top up to 5 mL for samples < 3 mL Top up to 10 mL for samples ≥ 3 mL | | |
| ŭ | Place the tube (without lid) into the magnet and incubate. | RT for 10 minutes | RT for 10 minutes | | |
| 9 | Carefully pipette* (do not pour) off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells. | Discard supernatant | Discard supernatant | | |
| 10 | Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times. | Top up to 2.5 mL | Top up to 5 mL for samples < 3 mL Top up to 10 mL for samples ≥ 3 mL | | |
| | Place the tube (without lid) into the magnet and incubate. | RT for 5 minutes | RT for 5 minutes | | |
| 11 | Repeat steps as indicated. | Steps 9 and 10 (total of 1 x 10-minute and 2 x 5-minute separations) | Steps 9 and 10 (total of 1 x 10-minute and 2 x 5-minute separations) | | |
| 12 | Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube. | Isolated cells are ready for use | Isolated cells are ready for use | | |

RT - room temperature (15 - 25°C)

^{*} Collect the entire supernatant, all at once, into a single pipette (e.g. for EasyEights™ 5 mL tube, use a 2 mL serological pipette [Catalog #38002]; for EasyEights™ 14 mL tube, use a 10 mL serological pipette [Catalog #38004]).



Directions for Use – Fully Automated RoboSep™ Protocol

See page 2 for Sample Preparation and Recommended Medium. Refer to Table 3 for detailed instructions regarding the RoboSep™ procedure.

Table 3. RoboSep™ Mouse CD11b Positive Selection Kit II Protocol for BONE MARROW

| STEP | INSTRUCTIONS | RoboSep™ (Catalog #21000) | | |
|------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|--|--|
| 1 | Prepare sample at the indicated cell concentration within the volume range. | 1 x 10^8 cells/mL 0.5 - 3 mL | | |
| 2 | Add Mouse FcR PolyBlock to sample. | 50 μL/mL of sample | | |
| 3 | Add sample to required tube. | 14 mL (17 x 95 mm) polystyrene round-bottom tube (e.g. Catalog #38008) | | |
| 4 | Prepare Selection Cocktail in the RoboSep™ Empty Vial provided. See Table 4 for required volumes. | Mix equal volumes of Component A and Component B (see Table 4). Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | | |
| | Incubate. | RT for 5 minutes | | |
| 5 | Select protocol. | Mouse CD11b Positive Selection II 18970 Bone marrow | | |
| 6 | Vortex RapidSpheres™. | 30 seconds | | |
| - | Load the carousel. | Follow on-screen prompts | | |
| | Start the protocol. | Press the green "Run" button | | |
| 8 | Unload the carousel when the run is complete. Remove the tube containing the isolated cells and resuspend in desired medium. Be sure to collect cells from the sides of the tube. | Isolated cells are ready for use | | |

Table 4. RoboSep™ Selection Cocktail Preparation

| START SAMPLE | COMPONENT A | COMPONENT B | SELECTION COCKTAIL TOTAL VOLUME |
|--------------|-------------|-------------|------------------------------------|
| 1 mL | 75 µL | 75 µL | 150 μL |
| 1.5 mL | 87.5 μL | 87.5 μL | 175 μL |
| 2 mL | 100 μL | 100 μL | 200 μL |
| 3 mL | 125 µL | 125 µL | 250 μL |
| 4 mL | 150 μL | 150 μL | 300 μL |

NOTE: RoboSep™ requires an excess of the Selection Cocktail to run properly (as indicated above).



Notes and Tips

ASSESSING PURITY


For purity assessment by flow cytometry, use the following fluorochrome-conjugated antibody clone:


Anti-Mouse CD11b Antibody, Clone M1/70 (Catalog #60001) at a concentration of 5 μg/mL

The following methods can also be used:

- Use a fluorochrome-conjugated secondary antibody, such as Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal (Catalog #60138).
- Add fluorochrome-conjugated Anti-Mouse CD11b Antibody, Clone M1/70 at a concentration of 0.5 µg/mL immediately after adding the cocktail. This method labels the positive cells in the entire sample.

Data





Starting with bone marrow samples, the CD11b+ cell content of the isolated fraction is typically 99.7 \pm 0.3% (mean \pm SD using the purple EasySepTM Magnet). In the above example, the purities of the start and final isolated fractions are 66.8% and 99.6%, respectively.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO WWW.STEMCELL.COM/COMPLIANCE.

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasyEights, EasySep, RapidSpheres, and RoboSep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.