

# EasySep™ Mouse F4/80 Positive Selection Kit



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

**For processing  $2 \times 10^9$  cells from spleen**

Catalog #100-0659

Positive Selection

Document #10000010445 | Version 01

## Description

Isolate highly purified F4/80+ cells from mouse splenocytes by positive selection.

- Fast, easy-to-use, and column-free
- Up to 92% purity
- Isolated cells are not fluorochrome-labeled

This kit targets F4/80+ cells for positive selection with antibodies recognizing the F4/80 cell surface marker. Desired cells are labeled with antibodies and magnetic particles and separated without columns using an EasySep™ magnet. Unwanted cells are simply poured off while desired cells remain in the tube. Isolated cells are immediately available for downstream applications such as flow cytometry, cell culture, and cell-based experiments.

NOTE: This is the Product Information Sheet (PIS) for isolating F4/80+ cells from mouse splenocytes. If isolating F4/80+ cells from mouse peritoneal lavage or lung tissue, refer to the applicable PIS, available at [www.stemcell.com](http://www.stemcell.com), or contact us to request a copy.

## Component Descriptions

COMPONENT NAME	COMPONENT #	QUANTITY	STORAGE	SHELF LIFE	FORMAT
EasySep™ Mouse F4/80 Positive Selection Component A	300-0251	1 x 0.3 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS and 0.1% BSA.
EasySep™ Mouse F4/80 Positive Selection Component B	300-0253	1 x 0.6 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of monoclonal antibodies in PBS with 5% HPCD.
EasySep™ Dextran RapidSpheres™ 50100	50100	2 x 1 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A suspension of magnetic particles in water.
Mouse FcR PolyBlock	300-0902	1 x 1.2 mL	Store at 2 - 8°C. Do not freeze.	Stable until expiry date (EXP) on label.	A combination of polyclonal antibodies and maltose in water with 5 µg/mL Triton X-100.

BSA - bovine serum albumin; HPCD - 2-hydroxypropyl-β-cyclodextrin; PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

## Sample Preparation

### SPLEEN

1. Harvest spleen into a conical tube with PBS containing 2% fetal bovine serum (FBS) or EasySep™ Buffer (Catalog #20144).
2. Place a 70 µm nylon mesh strainer (e.g. Catalog #27260) over a 50 mL conical tube (e.g. Catalog #38010) and push spleen tissue through the strainer with the rubber end of a syringe plunger to obtain a single-cell suspension.
3. Centrifuge at 300 x g for 10 minutes at room temperature with the brake on low. Carefully discard the supernatant.
4. Resuspend cells at  $1 \times 10^8$  nucleated cells/mL in recommended medium.

NOTE: Ammonium chloride treatment is not recommended when preparing the cells for separation.

### PERITONEAL LAVAGE OR LUNG TISSUE

If processing peritoneal lavage or lung tissue, refer to the applicable PIS, available at [www.stemcell.com](http://www.stemcell.com), or contact us to request a copy.



## Recommended Medium

EasySep™ Buffer (Catalog #20144) or PBS containing 2% FBS and 1 mM EDTA. Medium should be free of  $Ca^{++}$  and  $Mg^{++}$ .

## Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Tables 1 and 2 for detailed instructions regarding the EasySep™ procedure for each magnet.





**Table 1. EasySep™ Mouse F4/80 Positive Selection Kit Protocol for SPLEEN**

		EASYSEP™ MAGNETS	
STEP	INSTRUCTIONS	 <b>EasySep™</b> (Catalog #18000)	<b>“The Big Easy”</b> (Catalog #18001) 
1	Prepare sample at the indicated cell concentration within the volume range.	1 x 10 <sup>8</sup> cells/mL 0.5 - 2 mL	1 x 10 <sup>8</sup> cells/mL 1 - 8 mL
2	Add Mouse FcR PolyBlock to sample.	50 µL/mL of sample	50 µL/mL of sample
3	Add sample to required tube.	5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Catalog #38007)	14 mL (17 x 95 mm) polystyrene round-bottom tube (e.g. Catalog #38008)
4	Prepare Selection Cocktail in a tube. For each 1 mL of sample, prepare 45 µL of cocktail (15 µL of Component A + 30 µL of Component B).	Mix Component A and Component B at a 1:2 ratio. NOTE: Selection Cocktail must be prepared fresh before each use.	Mix Component A and Component B at a 1:2 ratio. NOTE: Selection Cocktail must be prepared fresh before each use.
	Incubate.	RT for 5 minutes	RT for 5 minutes
5	Add Selection Cocktail to sample. NOTE: Do not vortex cocktail.	45 µL/mL of sample	45 µL/mL of sample
	Mix and incubate.	RT for 5 minutes	RT for 5 minutes
6	Vortex RapidSpheres™. NOTE: Particles should appear evenly dispersed.	30 seconds	30 seconds
7	Add RapidSpheres™ to sample.	20 µL/mL of sample • For higher recovery, add 40 µL/mL of sample	20 µL/mL of sample • For higher recovery, add 40 µL/mL of sample
	Mix and incubate.	RT for 5 minutes	RT for 5 minutes
8	Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 2.5 mL	<ul style="list-style-type: none"> <li>• Top up to 5 mL for samples ≤ 3 mL</li> <li>• Top up to 10 mL for samples &gt; 3 mL</li> </ul>
	Place the tube (without lid) into the magnet and incubate.	RT for 3 minutes	RT for 5 minutes
9	Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells.	Discard supernatant	Discard supernatant
10	Repeat steps as indicated.	Steps 8 and 9, three more times (total of 4 x 3-minute separations)	Steps 8 and 9, three more times (total of 4 x 5-minute separations)
11	Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube or plate.	Isolated cells are ready for use	Isolated cells are ready for use

RT - room temperature (15 - 25°C)

\* Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.

Table 2. EasySep™ Mouse F4/80 Positive Selection Kit Protocol for SPLEEN

STEP	INSTRUCTIONS	EASYSEP™ MAGNETS		
		 <b>EasyPlate™</b> (Catalog #18102)	 <b>EasyEights™ (Catalog #18103)</b>	
			 <b>5 mL tube</b>	 <b>14 mL tube</b>
1	Prepare sample at the indicated cell concentration within the volume range.	1 x 10 <sup>8</sup> cells/mL 0.1 - 0.2 mL	1 x 10 <sup>8</sup> cells/mL 0.5 - 2 mL	1 x 10 <sup>8</sup> cells/mL 1 - 8 mL
2	Add Mouse FcR PolyBlock to sample.	50 µL/mL of sample	50 µL/mL of sample	50 µL/mL of sample
3	Add sample to required tube or plate.	Round-bottom, non-tissue culture-treated 96-well plate (e.g. Catalog #38018)	5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Catalog #38007)	14 mL (17 x 95 mm) polystyrene round-bottom tube (e.g. Catalog #38008)
4	Prepare Selection Cocktail in a tube. For each 1 mL of sample, prepare 45 µL of cocktail (15 µL of Component A + 30 µL of Component B).	Mix Component A and Component B at a 1:2 ratio. NOTE: Selection Cocktail must be prepared fresh before each use.	Mix Component A and Component B at a 1:2 ratio. NOTE: Selection Cocktail must be prepared fresh before each use.	Mix Component A and Component B at a 1:2 ratio. NOTE: Selection Cocktail must be prepared fresh before each use.
	Incubate.	RT for 5 minutes	RT for 5 minutes	RT for 5 minutes
5	Add Selection Cocktail to sample. NOTE: Do not vortex cocktail.	45 µL/mL of sample	45 µL/mL of sample	45 µL/mL of sample
	Mix and incubate.	RT for 5 minutes	RT for 5 minutes	RT for 5 minutes
6	Vortex RapidSpheres™. NOTE: Particles should appear evenly dispersed.	30 seconds	30 seconds	30 seconds
7	Add RapidSpheres™ to sample.	20 µL/mL of sample • For higher recovery, add 40 µL/mL of sample	60 µL/mL of sample • For higher recovery, add 80 µL/mL of sample	60 µL/mL of sample • For higher recovery, add 80 µL/mL of sample
	Mix and incubate.	RT for 5 minutes	RT for 5 minutes	RT for 5 minutes
8	Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 0.25 mL	Top up to 2.5 mL	• Top up to 5 mL for samples ≤ 3 mL • Top up to 10 mL for samples > 3 mL
	Place the tube or plate (without lid) into the magnet and incubate.	RT for 10 minutes	RT for 10 minutes	RT for 10 minutes
9	Carefully pipette* (do not pour) off the supernatant. Remove the tube or plate from the magnet; this tube or plate contains the isolated cells.	Discard supernatant	Discard supernatant	Discard supernatant
10	Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.	Top up to 0.25 mL	Top up to 2.5 mL	• Top up to 5 mL for samples ≤ 3 mL • Top up to 10 mL for samples > 3 mL
	Place the tube or plate (without lid) into the magnet and incubate.	RT for 5 minutes	RT for 5 minutes	RT for 5 minutes
11	Carefully pipette* (do not pour) off the supernatant. Remove the tube or plate from the magnet; this tube or plate contains the isolated cells.	Discard supernatant	Discard supernatant	Discard supernatant
12	Repeat steps as indicated.	Steps 10 and 11, two more times (total of 1 x 10-minute and 3 x 5-minute separations)	Steps 10 and 11, two more times (total of 1 x 10-minute and 3 x 5-minute separations)	Steps 10 and 11, two more times (total of 1 x 10-minute and 3 x 5-minute separations)
13	Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube or plate.	Isolated cells are ready for use	Isolated cells are ready for use	Isolated cells are ready for use

RT - room temperature (15 - 25°C)

\* Collect the entire supernatant, all at once, into a single pipette (e.g. for EasyEights™ 5 mL tube, use a 2 mL serological pipette [Catalog #38002]; for EasyEights™ 14 mL tube, use a 10 mL serological pipette [Catalog #38004]).

## Notes and Tips


### ASSESSING PURITY

For purity assessment by flow cytometry, use the following fluorochrome-conjugated antibody clones:

- Anti-Mouse F4/80 Antibody, Clone BM8 (Catalog #60027), and
- Anti-Mouse CD11b Antibody, Clone M1/70 (Catalog #100-0433), and
- Anti-Mouse CD45 Antibody, Clone 30-F11 (Catalog #60030)

NOTE: To exclude dead cells for purity assessment, use of the nuclear dye DRAQ7™ is recommended.

## Data



Starting with naïve mouse splenocytes, the F4/80+ cell content of the isolated fraction is typically  $89.8 \pm 2.9\%$  (mean  $\pm$  SD using the purple EasySep™ magnet). In the above example, the purities of the start and final isolated fractions are 5.4% and 90.6%, respectively.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED. FOR ADDITIONAL INFORMATION ON QUALITY AT STEMCELL, REFER TO [WWW.STEMCELL.COM/COMPLIANCE](http://WWW.STEMCELL.COM/COMPLIANCE).

Copyright © 2024 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasyEights, EasySep, EasyPlate, and RapidSpheres are trademarks of STEMCELL Technologies Canada Inc. DRAQ7 is a trademark of Biostatus Limited. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.