

Human Recombinant M-CSF (HEK293-expressed)

Macrophage colony-stimulating factor

Catalog #100-1723 100 µg

Catalog #100-1724 1000 µg

Product Description

Macrophage colony-stimulating factor (M-CSF) is a homodimeric glycoprotein growth factor that regulates proliferation and differentiation of myeloid hematopoietic progenitor cells to mononuclear phagocytic cell lineages, including monocytes, macrophages, and osteoclasts. M-CSF is a crucial factor for the development of tissue-resident macrophages in most tissues (Ginhoux & Jung). It is required for the maturation and activation of monocytes and macrophages, and regulates inflammatory responses in conjunction with other stimuli such as IFN-γ, LPS, and IL-4 (Murray et al.). M-CSF is also required for bone resorption by osteoclasts, and is involved in the development and regulation of placenta, mammary glands, and brain. M-CSF is produced by monocytes, fibroblasts, osteoclasts, stromal cells, endothelial cells, and tumor cells (Chockalingam & Ghosh).

M-CSF exerts its biological effects by signaling through a receptor tyrosine kinase (CSF-1R or M-CSF-R) encoded by the c-fms proto-oncogene (Hamilton). CSF-1R shares similar structural features with other growth factor receptors, including the stem cell factor (SCF) receptor, platelet-derived growth factor receptor (PDGF-R), and Flt3/Flk-2 receptor tyrosine kinase. Stimulation of the CSF-1R upon binding to M-CSF activates MAPK, PI3K, and PLCγ signaling pathways (Chockalingam & Ghosh). Human and mouse M-CSF sequences are highly conserved both at nucleotide and amino acid levels (80% homology; DeLamar et al.).

Product Information

Alternative Names:	Colony stimulating factor 1, CSF-1
Accession Number:	P09603
Amino Acid Sequence:	EEVSEYCSHM IGSGLHLSLQ RLIDSQMETS CQITFEFVDQ EQLKDPVCYL KKAFLLVQDI MEDTMRFRDN TPNAIAIVQL QELSLRLKSC FTKDYEEHDK ACVRTFYETP LQLEKVKNV FNETKNLLDK DWNIFSKNCN NSFAECSSQD VVTKPDCN
Predicted Molecular Mass:	18.4 kDa
Species:	Human
Product Formulation:	Lyophilized from sterile phosphate-buffered saline, pH 7.4, 5% Trehalose, 5% Mannitol, 0.01% Tween®-80.
Source:	HEK293
Purity:	≥ 82%

Specifications

Activity:	The specific activity is $\geq 1 \times 10^5$ units/mg ($EC_{50} \leq 6.3$ ng/mL), as determined by a cell proliferation assay using mouse M-NFS-60 cells.
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/ μ g protein.

Preparation and Storage

Stability and Storage:	Store at -20 to -80°C. Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.25 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data




Figure 1. Biological Activity and Molecular Mass of Human Recombinant M-CSF (HEK293-expressed)

(A) The biological activity of Human Recombinant M-CSF (HEK293-expressed) was tested by its ability to promote the proliferation of M-NFS-60 myelogenous leukemia lymphoblast cells. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC50 in the above example is ≤ 6.3 ng/mL.

(B) Human Recombinant M-CSF (HEK293-expressed) was resolved with SDS-PAGE under reducing conditions. Human Recombinant M-CSF has a predicted total molecular mass of 18.4 kDa. The recombinant protein migrates with an apparent molecular mass of 20 - 30 kDa.

Related Products

For a complete list of cytokines or peptide pools, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Chockalingam S & Ghosh SS. (2014) Macrophage colony-stimulating factor and cancer: a review. *Tumour Biol* 35(11): 10635–44.
- DeLamar JF et al. (1987) Nucleotide sequence of a cDNA encoding murine CSF-1 (Macrophage-CSF). *Nucleic Acids Res* 15(5): 2389–90.
- Ginhoux F & Jung S. (2014) Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nat Rev Immunol* 14(6): 392–404.
- Hamilton JA. (1997) CSF-1 signal transduction. *J Leukoc Biol* 62(2): 145–55.

PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2025 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.