

Anti-Human CD32 Antibody, Clone IV.3, FITC

Antibodies

Mouse monoclonal IgG2b antibody
against human CD32, FITC-conjugated

Catalog #60012FI.1
#60012FI

25 tests 20 µL/test
100 tests 20 µL/test



Scientists Helping Scientists™ | WWW.STEMCELL.COM

T. +1 604 877 0713 • TOLL-FREE T. 1 800 667 0322
ORDERS@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM
FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

FOR RESEARCH USE ONLY. NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES.

Product Description

The IV.3 antibody reacts with human CD32 (Fc_YRII), an ~40 kDa type 1 transmembrane glycoprotein that mediates several functions including phagocytosis, cytotoxicity, immunomodulation and platelet aggregation. CD32 is encoded by three genes (A, B, C) and at least 6 isoforms are generated via alternative mRNA splicing, i.e., IIa1, IIa2, IIb1, IIb2, IIb3 and IIc. All isoforms are expressed by monocytes/macrophages, placental trophoblasts and endothelial cells. In addition, the IIb isoform is expressed by B cells, and the IIa isoform by platelets, granulocytes and, weakly, by B cells. Isoform IIc is expressed by NK cells and neutrophils. CD32 binds weakly to the Fc region of monomeric IgG but more strongly to IgG aggregates and immune complexes. These interactions can result in non-specific labeling in antibody-based detection and cell separation experiments and the IV.3 antibody may be employed as a blocking antibody to reduce non-specific binding. The IV.3 antibody binds most strongly to the IIa isoforms of CD32.

Target Antigen Name:	CD32
Alternative Names:	FCR II, Fc _Y RII
Gene ID:	2212
Species Reactivity:	Human
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	IV.3
Isotype:	IgG2b
Immunogen:	K-562 human erythromyeloblastoid leukemia cell line
Conjugate:	FITC

Applications


Verified:	CellSep, FC
Reported:	FC
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human T Cell Enrichment Kit (Catalog #19051) and EasySep™ Human CD4+ T Cell Enrichment Kit (Catalog #19052).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; WB: Western blotting

Properties

Formulation:	Phosphate-buffered saline containing < 0.1% (w/v) sodium azide and < 0.1% (w/v) bovine serum albumin
Purification:	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to light. For product expiry date, please request a lot-specific Certificate of Analysis from techsupport@stemcell.com .
Directions for Use:	For flow cytometry the suggested use of this antibody is 20 µL per 1 x 10 ⁶ cells in 100 µL volume or per 100 µL of whole blood. It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD32 Antibody, Clone IV.3, FITC (filled histogram) or a mouse IgG2b, kappa FITC isotype control antibody (solid line histogram).

Related Products

PRODUCT NAME	CATALOG #	SIZE
Anti-Human CD32 Antibody, Clone IV.3	60012	100 µg
Anti-Human CD32 Antibody, Clone IV.3, FITC	60012FI.1	25 tests
Anti-Human CD32 Antibody, Clone IV.3, FITC	60012FI	100 tests

References

1. Looney RJ, et al. Human monocytes and U937 cells bear two distinct Fc receptors for IgG. *J Immunol* 136(5): 1641-47, 1986
2. Micklem KJ, et al. Different isoforms of human FcRII distinguished by CD32 antibodies. *J Immunol* 144(6): 22295-303, 1990 (FC, ICC, IHC, IP)
3. Tomiyama Y, et al. Response of human platelets to activating monoclonal antibodies: importance of Fc gamma RII (CD32) phenotype and level of expression. *Blood* 80(9): 2261-68, 1992
4. Ierino FL, et al. Mapping epitopes of human Fc gamma RII (CDw32) with monoclonal antibodies and recombinant receptors. *J Immunol* 150: 1794-803, 1993 (FC)
5. Schlossman SF, et al. Eds. *Leucocyte Typing V: Binding heterogeneity within the CD32 panel of mAB*. Oxford University Press, New York, pp. 832-35, 1995
6. Van Sorge NM, et al. FcgammaR polymorphisms: Implications for function, disease susceptibility and immunotherapy. *Tissue Antigens* 61(3): 189-202, 2003
7. Boruchov AM, et al. Activating and inhibitory IgG Fc receptors on human DCs mediate opposing functions. *J Clin Invest* 115(10): 2914-23, 2005 (FC)