

Anti-Human CD20 Antibody, Clone 2H7



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG2b antibody
against human, rhesus, cynomolgus
CD20, unconjugated

Catalog #60008

100 µg 0.5 mg/mL

Product Description

Antibody 2H7 reacts with CD20, an ~35 kDa non-glycosylated type 1 transmembrane protein in the MS4A protein family. The CD20 polypeptide transverse the cell membrane four times, with only a minor portion of the protein displayed on the cell surface. The epitope recognized by antibody 2H7 has been mapped to the amino acid sequence YNCEPANPSEKNSPST, located in the large extracellular loop of CD20. CD20 is expressed on pre-B cells, resting and activated B cells, some follicular dendritic cells, and a subset of T cells. Expression by B cells is lost upon their differentiation into plasma cells. By associating with several proteins, including the B cell receptor (CD79), MHC class I and II, CD53, CD81 and CD82, CD20 is involved in initiating intracellular signaling pathways that modulate the activation, proliferation and differentiation of B cells. It is thought that CD20 forms multi-subunit ion channels which regulate calcium ion flux across the plasma membrane. Activation of CD20 is accompanied by pronounced phosphorylation of the cytoplasmic domain of the ~33 kDa apo-protein, with the appearance of 35 - 37 kDa isoforms which associate with Src family kinases such as Fyn, Lck and Lyn.

Target Antigen Name:	CD20
Alternative Names:	B-lymphocyte antigen, B1, Bp35, Leu-16, MS4A1
Gene ID:	931
Species Reactivity:	Human, Rhesus, Cynomolgus, Baboon, Chimpanzee, Capuchin Monkey, Nancy Ma's Night Monkey, Pigtailed Macaque, Squirrel Monkey
Host Species:	Mouse
Clonality:	Monoclonal
Clone:	2H7
Isotype:	IgG2b, kappa
Immunogen:	Human tonsillar B cells
Conjugate:	Unconjugated

Applications


Verified:	FC
Reported:	CyTOF®, ELISA, FC, IHC, IP
Special Applications:	This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including EasySep™ Human Whole Blood CD20 Positive Selection Kit (Catalog #18685), EasySep™ Human CD19 Positive Selection Kit (Catalog #18054), and EasySep™ HLA Whole Blood Lymphoid Positive Selection Kit (Catalog #18684HLA).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	For flow cytometry the suggested use of this antibody is ≤ 2 µg per 1 × 10 ⁶ cells in 100 µL volume. It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD20 Antibody, Clone 2H7, followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (Catalog #60138FI) (filled histogram), or Mouse IgG2b, kappa Isotype Control Antibody, Clone MPC-11 (Catalog #60072), followed by Goat Anti-Mouse IgG (H+L) Antibody, Polyclonal, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Agrawal SM et al. (2013) Extracellular matrix metalloproteinase inducer shows active perivascular cuffs in multiple sclerosis. *Brain* 136(Pt 6): 1760–77. (ICC, IF)
2. Du J et al. (2008) Crystal structure of chimeric antibody C2H7 Fab in complex with a CD20 peptide. *Mol Immunol* 45(10): 2861–8.
3. Daubenberger CA et al. (2007) Flow cytometric analysis on cross-reactivity of human-specific CD monoclonal antibodies with splenocytes of Aotus nancymae, a non-human primate model for biomedical research. *Vet Immunol Immunopathol* 119(1-2): 14–20. (FC)
4. Teeling JL et al. (2006) The biological activity of human CD20 monoclonal antibodies is linked to unique epitopes on CD20. *J Immunol* 177(1): 362–71. (ELISA, FC)
5. Mack CL et al. (2004) Biliary atresia is associated with CD4+ Th1 cell-mediated portal tract inflammation. *Pediatr Res* 56(1): 79–87. (IHC)
6. Cragg MS et al. (2003) Complement-mediated lysis by anti-CD20 mAb correlates with segregation into lipid rafts. *Blood* 101(3): 1045–52. (FA, FC)
7. Mason D. (2002) *Leucocyte Typing VII*. D. Mason, ed., New York: Oxford University Press.
8. Barclay AN et al. (Eds.). (1997) *The Leucocyte Antigen FactsBook*, Second Edition. New York: Academic Press.
9. Kanzaki M et al. (1997) Activation of a calcium-permeable cation channel CD20 expressed in Balb/c 3T3 cells by insulin-like growth factor-I. *J Biol Chem* 272(8): 4964–9.
10. Szöllösi J et al. (1996) Supramolecular complexes of MHC class I, MHC class II, CD20, and tetraspan molecules (CD53, CD81, and CD82) at the surface of a B cell line JY. *J Immunol* 157(7): 2939–46.
11. Tedder TF & Engel P. (1994) CD20: a regulator of cell-cycle progression of B lymphocytes. *Immunol Today* 15(9): 450–4.
12. Hultin LE et al. (1993) CD20 (pan-B cell) antigen is expressed at a low level on a subpopulation of human T lymphocytes. *Cytometry* 14(2): 196–204.
13. Liu AY et al. (1987) Production of a mouse-human chimeric monoclonal antibody to CD20 with potent Fc-dependent biologic activity. *J Immunol* 139(10): 3521–6. (FA)
14. McMichael AJ et al. (Eds.). (1987) *Leucocyte Typing III: White Cell Differentiation Antigens*. New York: Oxford University Press.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.