Anti-Mouse Ly-6G Antibody, Clone 1A8

Antibodies

Rat monoclonal IgG2a antibody against mouse Ly-6G, unconjugated

Catalog #60031 500 μg 0.5 mg/mL



Scientists Helping Scientists[™] | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

The 1A8 antibody reacts with Ly-6G, a 21 - 25 kDa GPI-anchored protein, which together with the structurally related Ly-6C protein comprises the granulocyte receptor-1 antigen (Gr-1). Gr-1 is expressed on monocytes, neutrophils and subsets of macrophages, plasmacytoid dendritic cells and T cells. Monocytes in the bone marrow transiently express Gr-1 during development and the expression level is strongly correlated with granulocyte differentiation and maturation. In the periphery, Gr-1 is found predominantly on neutrophils and is a useful marker for these cells. The 1A8 antibody binds specifically to Ly-6G, whereas another commonly used antibody, clone RB6-8C5, binds to both Ly-6G and Ly-6C. It has been reported that the 1A8 antibody detects Ly-6G-expressing granulocytes in peripheral blood, whereas the RB6-8C5 antibody also binds to Ly-6C-expressing lymphocytes, monocytes and dendritic cells.

Target Antigen Name: Ly-6G

Alternative Names: Gr-1, GR1, Ly6G

Gene ID: 546644

Species Reactivity: Mouse

Host Species: Rat (LEW)

Clonality: Monoclonal

Clone: 1A8

Isotype: IgG2a, kappa

Immunogen: Mouse Ly-6G-transfected EL-4J cell line

Conjugate: Unconjugated

Applications

Verified: FC

Reported: FA, FC, IP, IHC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Mouse Neutrophil Enrichment Kit (Catalog #19762).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Purification: The antibody was purified by affinity chromatography.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact

techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is ≤ 0.25 µg per 1 x 10^6 cells in 100 µL volume. It is


recommended that the antibody be titrated for optimal performance for each application.


Antibodies

Anti-Mouse Ly-6G Antibody, Clone 1A8



Data





(A) Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse Ly-6G Antibody, Clone 1A8, followed by a mouse anti-rat IgG2a antibody, FITC (filled histogram), or Rat IgG2a, kappa Isotype Control Antibody, Clone RTK2758 (Catalog #60076), followed by a mouse anti-rat IgG2a antibody, FITC (solid line histogram).

(B) Immunohistochemical analysis of formalin-fixed, paraffin-embedded mouse uterine tissue labeled with Anti-Mouse Ly-6G Antibody, Clone 1A8, followed by anti-mouse, HRP. DAB substrate was used for visualization.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- 1. Rose S et al. (2012) A novel Ly6C/Ly6G-based strategy to analyze the mouse splenic myeloid compartment. Cytometry A 81(4): 343–50. (FC)
- 2. Fujita M et al. (2011) COX-2 blockade suppresses gliomagenesis by inhibiting myeloid-derived suppressor cells. Cancer Res 71(7): 2664-74.
- 3. Guiducci C et al. (2010) Autoimmune skin inflammation is dependent on plasmacytoid dendritic cell activation by nucleic acids via TLR7 and TLR9. J Exp Med 207(13): 2931–42.
- 4. Tadagavadi RK & Reeves WB. (2010) Endogenous IL-10 attenuates cisplatin nephrotoxicity: role of dendritic cells. J Immunol 185(8): 4904–11.
- 5. Daley JM et al. (2008) Use of Ly6G-specific monoclonal antibody to deplete neutrophils in mice. J Leukoc Biol 83(1): 64–70.
- 6. van Leeuwen M et al. (2008) Accumulation of myeloperoxidase-positive neutrophils in atherosclerotic lesions in LDLR-/- mice. Arterioscler Thromb Vasc Biol 28(1): 84–9. (IHC)
- 7. Dietlin TA et al. (2007) Mycobacteria-induced Gr-1+ subsets from distinct myeloid lineages have opposite effects on T cell expansion. J Leukoc Biol 81(5): 1205–12. (FC)
- 8. Fleming TJ & Malek TR. (1994) Multiple glycosylphosphatidylinositol-anchored Ly-6 molecules and transmembrane Ly-6E mediate inhibition of IL-2 production. J Immunol 153(5): 1955–62. (FA)
- 9. Fleming TJ et al. (1993) Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. J Immunol 151(5): 2399–408. (IP)
- 10. Hestdal K et al. (1991) Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. J Immunol 147(1): 22-8.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.