

Positive Selection

Catalog #18954

For processing 2 x 10<sup>9</sup> cells



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Document #28032 | Version 2\_0\_1

### Description

Isolate highly purified CD19+ cells from single-cell suspensions of mouse splenocytes or other tissues by immunomagnetic positive selection.

- Fast and easy-to-use
- · Up to 99% purity
- · No columns required
- · Isolated cells are not fluorochrome labeled

This kit targets CD19+ cells for positive selection with antibodies recognizing the CD19 surface marker. Desired cells are labeled with antibodies and magnetic particles, and separated without columns using an EasySep<sup>TM</sup> magnet. Unwanted cells are simply poured off, while desired cells remain in the tube. Isolated cells are immediately available for downstream applications such as flow cytometry, culture cell-based experiments.

### Component Descriptions

| COMPONENT NAME   | COMPONENT # | QUANTITY   | STORAGE                             | SHELF LIFE                               | FORMAT  |
|--|-------------|------------|-------------------------------------|--|---|
| EasySep™ Mouse CD19 Positive<br>Selection Kit II Component A | 18954CA     | 1 x 0.5 mL | Store at 2 - 8°C. Do not freeze.    | Stable until expiry date (EXP) on label. | A combination of monoclonal antibodies in PBS with 0.1% BSA and 10% HPCD. |
| EasySep™ Mouse CD19 Positive<br>Selection Kit II Component B | 18954CB     | 1 x 0.5 mL | Store at 2 - 8°C. Do not freeze.    | Stable until expiry date (EXP) on label. | A combination of monoclonal antibodies in PBS with 0.1% BSA and 10% HPCD. |
| EasySep™ Dextran<br>RapidSpheres™ 50100                      | 50100       | 1 x 1 mL   | Store at 2 - 8°C.<br>Do not freeze. | Stable until expiry date (EXP) on label. | A suspension of magnetic particles in water.                              |
| RoboSep™ Empty Vial  | 27401       | 1          | Not applicable                      | Not applicable                           | Not applicable  |

BSA - bovine serum albumin; HPCD - 2-hydroxypropyl-β-cyclodextrin; PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) and should be stored according to their storage conditions upon receipt.

### Additional Reagent Stability Information

| REAGENT NAME  | STORAGE                          | SHELF LIFE  |
|---|----------------------------------|---|
| Selection Cocktail (combined Component A + Component B) | Store at 2 - 8°C. Do not freeze. | Stable for up to 4 weeks. Do not exceed expiry date (EXP) of individual components. |

#### Sample Preparation

**SPLEEN** 

Disrupt spleen in PBS or Hanks' Balanced Salt Solution containing 2% fetal bovine serum (FBS). Remove clumps and debris by passing cell suspension through a 70 µm mesh nylon strainer. Centrifuge at 300 x g for 10 minutes and resuspend at 1 x 10^8 nucleated cells/mL in recommended medium.

Ammonium chloride treatment is not recommended when preparing the cells for separation.

### Recommended Medium

EasySep™ Buffer (Catalog #20144), RoboSep™ Buffer (Catalog #20104), or PBS containing 2% FBS and 1 mM EDTA. Medium should be free of Ca++ and Mg++.





### Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Tables 1 and 2 for detailed instructions regarding the EasySep™ procedure for each magnet.

#### Table 1. EasySep™ Mouse CD19 Positive Selection Kit II Protocol

| <b>1</b> |  | EASYSEP™ MAGNETS  |   |  |
|----------|--|---|---|--|
| STEP     | INSTRUCTIONS   | EasySep™ (Catalog #18000)   | "The Big Easy"<br>(Catalog #18001)  |  |
|          | Prepare sample at the indicated cell concentration within the volume range.  | 1 x 10^8 cells/mL<br>0.25 - 2 mL  | 1 x 10^8 cells/mL<br>0.5 - 8 mL   |  |
|          | Add sample to required tube.   | 5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352058)                              | 14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)                            |  |
| 2        | Prepare Selection Cocktail in a tube. For each 1 mL of sample make 50 μL of cocktail (25 μL of Component A + 25 μL of Component B).  | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. |  |
|          | Incubate.  | RT for 5 minutes  | RT for 5 minutes  |  |
|          | Add Selection Cocktail to sample.  | 50 μL/mL of sample  | 50 μL/mL of sample  |  |
| 3        | Mix and incubate.  | RT for 3 minutes  | RT for 3 minutes  |  |
| 4        | Vortex RapidSpheres™.  | 30 seconds  | 30 seconds  |  |
|          | Add RapidSpheres™ to sample.   | 75 μL/mL of sample  | 75 μL/mL of sample  |  |
| 5        | Mix and incubate.  | RT for 3 minutes  | RT for 3 minutes  |  |
| 6        | Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.  | Top up to 2.5 mL  | <ul> <li>Top up to 3 mL for samples &lt; 2.5 mL</li> <li>Top up to 10 mL for samples ≥ 2.5 mL</li> </ul>    |  |
|          | Place the tube (without lid) into the magnet and incubate.   | RT for 3 minutes  | RT for 3 minutes  |  |
| 7        | Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells. | Discard supernatant   | Discard supernatant   |  |
| 8        | Repeat steps as indicated.   | Steps 6 and 7 (total of 2 x 3-minute separations)   | Steps 6 and 7<br>(total of 2 x 3-minute separations)  |  |
| 9        | Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube.  | Isolated cells are ready for use  | Isolated cells are ready for use  |  |

RT - room temperature (15 - 25°C)

<sup>\*</sup> Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.





Table 2. EasySep™ Mouse CD19 Positive Selection Kit II Protocol

|      |   |   | EASYSEF  | MAGNETS   |  |  |
|------|---|---|--|---|--|--|
|      | INSTRUCTIONS  | EasyEights™ (Catalog #18103)  |  |   |  |  |
| STEP |   |   | 5 mL tube  | 14 mL tube  |  |  |
| 1    | Prepare sample at the indicated cell concentration within the volume range.   |   | 1 x 10^8 cells/mL<br>0.5 - 2 mL  | 1 x 10^8 cells/mL<br>0.5 - 8 mL   |  |  |
|      | Add sample to required tube.  | 5 mL (12  | x 75 mm) polystyrene round-bottom tube<br>(e.g. Corning Catalog #352058) | 14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)                            |  |  |
| 2    | Prepare Selection Cocktail in a tube. For each 1 mL of sample make 50 μL of cocktail (25 μL of Component A + 25 μL of Component B). | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. |  | Mix equal volumes of Component A and Component B. Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. |  |  |
|      | Incubate.   |   | RT for 5 minutes   | RT for 5 minutes  |  |  |
| •    | Add Selection Cocktail to sample.   |   | 50 μL/mL of sample   | 50 μL/mL of sample  |  |  |
| 3    | Mix and incubate.   |   | RT for 3 minutes   | RT for 3 minutes  |  |  |
| 4    | Vortex RapidSpheres™.   | 30 seconds  |  | 30 seconds  |  |  |
|      | Add RapidSpheres™ to sample.  |   | 75 μL/mL of sample   | 75 μL/mL of sample  |  |  |
| 5    | Mix and incubate.   |   | RT for 3 minutes   | RT for 3 minutes  |  |  |
| 6    | Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.                   | Top up to 2.5 mL  |  | <ul> <li>Top up to 3 mL for samples &lt; 2.5 mL</li> <li>Top up to 10 mL for samples ≥ 2.5 mL</li> </ul>    |  |  |
|      | Place the tube (without lid) into the magnet and incubate.  | RT for 10 minutes   |  | RT for 10 minutes   |  |  |
| 7    | Carefully pipette** (do not pour) off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells.      | Discard supernatant   |  | Discard supernatant   |  |  |
| 8    | Add recommended medium to top up sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times.                   | Top up to 2.5 mL  |  | <ul> <li>Top up to 3 mL for samples &lt; 2.5 mL</li> <li>Top up to 10 mL for samples ≥ 2.5 mL</li> </ul>    |  |  |
|      | Place the tube (without lid) into the magnet and incubate.  | RT for 5 minutes  |  | RT for 5 minutes  |  |  |
| 9    | Carefully pipette** (do not pour) off the supernatant. Remove the tube from the magnet; this tube contains the isolated cells.      | Discard supernatant Discard supernatant   |  | Discard supernatant   |  |  |
| 10   | Resuspend cells in desired medium. Be sure to collect cells from the sides of the tube.   |   | Isolated cells are ready for use   | Isolated cells are ready for use  |  |  |

RT - room temperature (15 - 25°C)

<sup>\*\*</sup> Collect the entire supernatant, all at once, into a single pipette (e.g. for the EasyEights™ 5 mL tube use a 2 mL serological pipette and for the EasyEights™ 14 mL tube use a 10 mL serological pipette).





### Directions for Use – Fully Automated RoboSep™ Protocol

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 3 for detailed instructions regarding the RoboSep™ procedure.

#### Table 3. RoboSep™ Mouse CD19 Positive Selection Kit Protocol

| STEP | INSTRUCTIONS  | RoboSep™<br>(Catalog #20000 and #21000)  |  |  |
|------|---|--|--|--|
| 1    | Prepare sample at the indicated cell concentration within the volume range.   | 1 x 10^8 cells/mL<br>0.5 - 8 mL  |  |  |
|      | Add sample to required tube.  | 14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057)   |  |  |
| 2    | Prepare Selection Cocktail in the RoboSep™ Empty Vial provided. See Table 4 for required volumes.   | Mix equal volumes of Component A and Component B (see Table 4).  Prepared cocktail is stable at 2 - 8°C for up to 4 weeks. |  |  |
|      | Incubate.   | RT for 5 minutes   |  |  |
| 3    | Select protocol.  | Mouse CD19 Positive Selection II 18954v2   |  |  |
| 4    | Vortex RapidSpheres™.   | 30 seconds   |  |  |
| 5    | Load the carousel.  | Follow on-screen prompts   |  |  |
| 5    | Start the protocol.   | Press the green "Run" button   |  |  |
| 6    | Unload the carousel when the run is complete. Remove the tube containing the isolated cells and resuspend in desired medium. Be sure to collect cells from the sides of the tube. | Isolated cells are ready for use   |  |  |

RT - room temperature (15 - 25°C)

Table 4. RoboSep™ Selection Cocktail Preparation

| START SAMPLE | COMPONENT A | COMPONENT B | SELECTION COCKTAIL<br>TOTAL VOLUME |
|--------------|-------------|-------------|------------------------------------|
| 0.5 mL       | 62.5 μL     | 62.5 μL     | 125 μL                             |
| 1 mL         | 75 μL       | 75 μL       | 150 μL                             |
| 1.5 mL       | 87.5 μL     | 87.5 μL     | 175 μL                             |
| 2 mL         | 100 μL      | 100 μL      | 200 μL                             |
| 3 mL         | 125 µL      | 125 µL      | 250 μL                             |
| 4 mL         | 150 μL      | 150 µL      | 300 μL                             |
| 5 mL         | 175 µL      | 175 µL      | 350 μL                             |
| 6 mL         | 200 μL      | 200 μL      | 400 μL                             |
| 7 mL         | 225 μL      | 225 µL      | 450 μL                             |
| 8 mL         | 250 μL      | 250 μL      | 500 μL                             |

Note: RoboSep™ requires an extra 100 µL of the Selection Cocktail to run properly (compared to manual protocols).






### Notes and Tips


ASSESSING PURITY

For purity assessment by flow cytometry use the following fluorochrome-conjugated antibody clone:

Anti-mouse CD19 antibody, clone 1D3 (partially blocked)

#### Data





Starting with mouse splenocytes, the CD19+ cell content of the isolated fraction is typically 98.1 ± 0.6% (mean ± SD using the purple EasySep™ Magnet).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasyEights, EasySep, RapidSpheres, and RoboSep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.