



Negative Selection

Catalog #19856

**EasySep™ Mouse
Hematopoietic Progenitor
Cell Isolation Kit**

For processing 1 x 10⁹ cells



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Document #28035 | Version 1_0_2

Description

Isolate untouched and highly purified progenitor cells from mouse splenocytes by immunomagnetic negative selection. When using single-cell suspensions from other tissue types, this kit may require optimization.

- Fast and easy-to-use
- Up to 84% purity
- No columns required
- Untouched, viable cells

This kit targets non-progenitor cells for removal with biotinylated antibodies recognizing specific cell surface markers. Unwanted cells are labeled with biotinylated antibodies and magnetic particles, and separated without columns using an EasySep™ magnet. Desired cells are simply poured off into a new tube. Isolated cells are immediately available for downstream applications such as flow cytometry, culture and cell-based experiments.

Component Descriptions

| COMPONENT NAME | COMPONENT # | QUANTITY | STORAGE | SHELF LIFE | FORMAT |
|---|-------------|------------|-------------------------------------|--|---|
| EasySep™ Mouse Hematopoietic Progenitor Cell Isolation Cocktail | 19856C | 1 x 0.5 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A combination of monoclonal antibodies in PBS and 0.1% BSA. |
| EasySep™ Streptavidin RapidSpheres™ 50001 | 50001 | 1 x 1 mL | Store at 2 - 8°C. Do not freeze. | Stable until expiry date (EXP) on label. | A suspension of magnetic particles in PBS. |
| Normal Rat Serum | 13551 | 1 x 2 mL | Store at -20°C. | Stable until expiry date (EXP) on label. | Mycoplasma-free normal rat serum. |

BSA - bovine serum albumin; PBS - phosphate-buffered saline

Components may be shipped at room temperature (15 - 25°C) but should be stored as indicated above.

Additional Reagent Stability Information

| REAGENT NAME | STORAGE | SHELF LIFE |
|---------------------------|-------------------|---|
| Normal Rat Serum (in-use) | Store at 2 - 8°C. | Stable for at least 2 months. Do not exceed expiry date (EXP) on label. |

Sample Preparation

BONE MARROW

Flush bone marrow cells from femur and tibia into recommended medium using a syringe equipped with a 23 gauge needle. Disperse clumps by gently passing the cell suspension through the syringe several times. Alternatively, crush bones using a mortar and pestle. Remove remaining clumps and debris by passing cell suspension through a 70 µm mesh nylon strainer. Centrifuge at 300 x g for 10 minutes and resuspend cells at 1 x 10⁸ cells/mL in recommended medium.

Recommended Medium

EasySep™ Buffer (Catalog #20144), RoboSep™ Buffer (Catalog #20104), or PBS (Catalog #37350) containing 2% fetal bovine serum (FBS) and 1 mM EDTA. Medium should be free of Ca++, Mg++, and biotin.

Directions for Use – Manual EasySep™ Protocols

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 1 for detailed instructions regarding the EasySep™ procedure for each magnet.

Table 1. EasySep™ Mouse Hematopoietic Progenitor Cell Isolation Kit Protocol

| | | EASYSEP™ MAGNETS | |
|------|---|--|---|
| STEP | INSTRUCTIONS | EasySep™ (Catalog #18000) | “The Big Easy” (Catalog #18001) |
| 1 | Prepare sample at the indicated cell concentration within the volume range. | 1 x 10 ⁸ cells/mL 0.5 - 2 mL | 1 x 10 ⁸ cells/mL 0.5 - 8 mL |
| 2 | Add Rat Serum to sample. | 50 µL/mL of sample | 50 µL/mL of sample |
| 3 | Add sample to required tube. | 5 mL (12 x 75 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352058) | 14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057) |
| 4 | Add Isolation Cocktail to sample. | 50 µL/mL of sample | 50 µL/mL of sample |
| | Mix and incubate. | 2 - 8°C for 15 minutes | 2 - 8°C for 15 minutes |
| 5 | Vortex RapidSpheres™. | 30 seconds | 30 seconds |
| 6 | Add RapidSpheres™ to sample. | 75 µL/mL of sample | 75 µL/mL of sample |
| | Mix and incubate. | 2 - 8°C for 10 minutes | 2 - 8°C for 10 minutes |
| 7 | Add recommended medium to top up the sample to the indicated volume. Mix by gently pipetting up and down 2 - 3 times. | Top up to 2.5 mL | <ul style="list-style-type: none"> Top up to 5 mL for samples ≤ 4 mL Top up to 10 mL for samples > 4 mL |
| | Place the tube (without lid) into the magnet and incubate. | RT for 3 minutes | RT for 3 minutes |
| 8 | Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the enriched cell suspension into a new tube. | Use a new 14 mL tube | <ul style="list-style-type: none"> Use a new 14 mL tube for start samples ≤ 4 mL Use a new 50 mL tube for start samples > 4 mL |
| 9 | Remove the tube from the magnet and add recommended medium to the indicated volume. Mix by gently pipetting up and down 5 - 6 times. | Top up to 2.5 mL | <ul style="list-style-type: none"> Top up to 5 mL for samples ≤ 4 mL Top up to 10 mL for samples > 4 mL |
| 10 | Place the tube (without lid) into the magnet and incubate. | RT for 3 minutes | RT for 3 minutes |
| 11 | Pick up the magnet, and in one continuous motion invert the magnet and tube,* pouring off the enriched cell suspension. | Combine with first poured-off fraction from step 8 Isolated cells are now ready for use | Combine with first poured-off fraction from step 8 Isolated cells are now ready for use |

RT - room temperature (15 - 25°C)

* Leave the magnet and tube inverted for 2 - 3 seconds, then return upright. Do not shake or blot off any drops that may remain hanging from the mouth of the tube.

Directions for Use – Fully Automated RoboSep™ Protocol

See page 1 for Sample Preparation and Recommended Medium. Refer to Table 2 for detailed instructions regarding the RoboSep™ procedure.

Table 2. RoboSep™ Mouse Hematopoietic Progenitor Cell Isolation Kit Protocol

| STEP | INSTRUCTIONS | RoboSep™ (Catalog #20000 and #21000) |
|------|---|---|
| 1 | Prepare sample at the indicated cell concentration within the volume range. | 1 x 10 ⁸ cells/mL 0.5 - 8 mL |
| 2 | Add Rat Serum to sample. | 50 µL/mL of sample |
| 3 | Add sample to required tube. | 14 mL (17 x 100 mm) polystyrene round-bottom tube (e.g. Corning Catalog #352057) |
| 4 | Select protocol. | Mouse Hematopoietic Progenitor Cell Isolation 19856 |
| 5 | Vortex RapidSpheres™. | 30 seconds |
| 6 | Load the carousel. | Follow on-screen prompts |
| | Start the protocol. | Press the green “Run” button |
| 7 | Unload the carousel when the run is complete. | Isolated cells are now ready for use |

Notes and Tips

ASSESSING PURITY

The first step for isolation of mouse hematopoietic stem and progenitor cells (HSPCs) from bone marrow consists of removing mature cells that express 'lineage' (Lin) antigens specific to terminally differentiated blood cells. Lineage antigens are absent or weakly expressed on HSPCs. Lineage antigens include CD3, CD11b, CD19, CD45R (B220), Ly6G/C (Gr-1), and TER119. In many mouse strains, HSPCs are positive for Sca1 (Ly-6A/E) and c-Kit (the receptor for SCF, also known as CD117) and referred to as LSK (Lin-Sca1+c-Kit+).^{1,2,3,4} LSK cells, which make up < 0.1% of nucleated BM cells, contain most repopulating stem cells but are depleted of more mature erythroid, myeloid and megakaryocyte cells including most colony-forming units (CFUs) which are Lin-Sca1-/^{low}c-Kit+.⁵ Mouse HSPCs are heterogeneous for other antigens such as CD34 and Thy1.


For purity assessment of these subsets of progenitor cells by flow cytometry use the following fluorochrome-conjugated antibody clones:

- Anti-Mouse CD117 Antibody, Clone 2B8 (Catalog #60025), and
- Anti-Mouse Sca1 Antibody, Clone E13-161.7 (Catalog #60032), and
- Anti-mouse lineage antibodies (see below)

For lineage antigen labeling use the following fluorochrome-conjugated antibody clones:

- Anti-Mouse CD3e Antibody, Clone 145-2C11 (Catalog #60015), and
- Anti-Mouse CD11b Antibody, Clone M1/70 (Catalog #60001), and
- Anti-Mouse CD19 Antibody, Clone 6D5 (Catalog #60006), and
- Anti-Mouse CD45R Antibody, Clone RA3-6B2 (Catalog #60019), and
- Anti-Mouse Gr-1 Antibody, Clone RB6-8C5 (Catalog #60028), and
- Anti-Mouse TER119 Antibody, Clone TER-119 (Catalog #60033)

Data



Starting with a mouse bone marrow cell suspension, the lineage antigen-negative cell content of the isolated fraction typically ranges from 60 - 84%.

References

1. Spangrude GJ et al. (1988) Purification and characterization of mouse hematopoietic stem cells. *Science* 241: 58–62.
2. Uchida N & Weissman IL. (1992) Searching for hematopoietic stem cells: evidence that Thy-1.1lo Lin- Sca-1+ cells are the only stem cells in C57BL/Ka-Thy-1.1 bone marrow. *J Exp Med* 175(1): 175–84.
3. Okada S et al. (1992) In vivo and in vitro stem cell function of c-kit- and Sca-1-positive murine hematopoietic cells. *Blood* 80(12): 3044–50.
4. Osawa M et al. (1996) In vivo self-renewal of c-Kit+ Sca-1+ Lin(low/-) hemopoietic stem cells. *J Immunol* 156(9): 3207–14.
5. Akashi K et al. (2000) A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 404(6774): 193–7.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2015 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, EasySep, RapidSpheres, and RoboSep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.