

Small Molecules

(-)-Indolactam V

Protein kinase C (PKC) activator

Catalog # 72312
72314

300 µg
1 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM


FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

(-)-Indolactam V is an indole alkaloid compound that activates protein kinase C (PKC). It binds to the α , β , γ , δ , ϵ , and η isozymes of PKC with K_i values of 11, 6, 19, 8, 22, and 16 nM respectively (Kazanietz et al.; Masuda et al.).

Molecular Name:	(-)-Indolactam V
Alternative Names:	Not applicable
CAS Number:	90365-57-4
Chemical Formula:	$C_{17}H_{23}N_3O_2$
Molecular Weight:	301.4 g/mol
Purity:	≥ 97%
Chemical Name:	(2S,5S)-1,2,4,5,6,8-hexahydro-5-(hydroxymethyl)-1-methyl-2-(1-methylethyl)-3H-pyrrolo[4,3,2-gh]-1,4-benzodiazonin-3-one

Structure:



Properties

Physical Appearance:	White to off-white solid
Storage:	Product stable at 2 - 8°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	· DMSO ≤ 30 mM · Absolute ethanol ≤ 15 mM For example, to prepare a 5 mM stock solution in DMSO, resuspend 300 µg in 199 µL of fresh DMSO. Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application. Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Promotes differentiation to human and mouse pancreatic precursors from pluripotent stem cell-derived definitive endoderm (Borowiak et al.; Chen et al.; Thatava et al.).

References

- Borowiak M et al. (2009) Small molecules efficiently direct endodermal differentiation of mouse and human embryonic stem cells. *Cell Stem Cell* 4(4): 348–58.
- Chen S et al. (2009) A small molecule that directs differentiation of human ESCs into the pancreatic lineage. *Nat Chem Biol* 5(4): 258–65.
- Kazanietz MG et al. (1993) Characterization of ligand and substrate specificity for the calcium-dependent and calcium-independent protein kinase C isozymes. *Mol Pharmacol* 44(2): 298–307.
- Masuda A et al. (2002) Binding selectivity of conformationally restricted analogues of (-)-indolactam-V to the C1 domains of protein kinase C isozymes. *Biosci Biotechnol Biochem* 66(7): 1615–7.
- Thatava T et al. (2011) Indolactam V/GLP-1-mediated differentiation of human iPS cells into glucose-responsive insulin-secreting progeny. *Gene Ther* 18(3): 283–93.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.