Anti-Mouse CD45R Antibody, Clone RA3-6B2, PerCP-Cv5.5

Antibodies

Rat monoclonal IgG2a antibody against human, mouse, cat CD45R (B220), PerCP-Cy5.5-conjugated

Catalog #60019PS #60019PS.1

100 μg25 μg0.2 mg/mL



Scientists Helping Scientists[™] | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

The RA3-6B2 antibody reacts with an epitope on the extracellular domain of the exon A-restricted isoform of mouse CD45. CD45 is a member of the protein tyrosine phosphatase family and functions in a number of immunoregulatory processes, including cell activation, growth, differentiation, and oncogenic transformation. The large cytoplasmic portion of CD45 contains two tyrosine phosphatase domains (one of which is enzymatically active) that are involved in modulating the function of Src kinases such as Lck and Fyn. Several isoforms of CD45 have been identified, the expression of which differs according to cell type and functional status. Alternative splicing of three exons encoding the extracellular RA, RB, and RC polypeptide sequences gives rise to up to 8 isoforms with molecular masses in the range of 200 - 220 kDa. The exon A-restricted isoform of mouse CD45, termed CD45R, contains the RA, RB, and RC regions and is the largest of the isoforms. CD45R is expressed by the B cell lineage from early pro-B to mature B cells, with its expression downregulated during differentiation to plasma cells. CD45R is also expressed on subsets of T, NK, dendritic, and lymphokine-activated killer (LAK) cells.

Target Antigen Name: CD45R (B220)

Alternative Names: B220, Protein tyrosine phosphatase receptor type C, PTPRC

Gene ID: 19264/5788

Species Reactivity: Human, Mouse, Cat

Host Species: Rat

Clonality:MonoclonalClone:RA3-6B2Isotype:IgG2a, kappa

Immunogen: Abelson murine leukemia virus-induced pre-B tumor cells

Conjugate: PerCP-Cy5.5 (Peridinin chlorophyll protein complex-Cyanine5.5)

Applications

Verified: FC Reported: FC

Special Applications: This antibody clone has been verified for purity assessments of cells isolated with EasySep™ kits, including

EasySep™ Mouse B Cell Isolation Kit (Catalog #19854).

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin

Purification: The antibody was purified by affinity chromatography and conjugated with PerCP-Cy5.5 under optimal

conditions. The solution is free of unconjugated PerCP-Cy5.5.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry, the suggested use of this antibody is ≤ 0.25 µg per 1 x 10⁶ cells in 100 µL. It is


recommended that the antibody be titrated for optimal performance for each application.

Anti-Mouse CD45R Antibody, Clone RA3-6B2, PerCP-Cy5.5

Antibodies



Data



Flow cytometry analysis of C57BL/6 mouse splenocytes labeled with Anti-Mouse CD45R Antibody, Clone RA3-6B2, PerCP-Cy5.5 (filled histogram) or a rat IgG2a, kappa isotype control antibody, PerCP-Cy5.5 (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

- 1. Masahata K et al. (2014) Generation of colonic IgA-secreting cells in the caecal patch. Nat Commun 5: 3704. (FC, IF, IHC)
- 2. Wang C et al. (2014) Lymphatic-targeted cationic liposomes: A robust vaccine adjuvant for promoting long-term immunological memory. Vaccine 32(42): 5475–83. (IF, IHC)
- 3. Nishino T, et al. (2012) Involvement of lymphocyte infiltration in the progression of mouse peritoneal fibrosis model. Ren Fail 34(6): 760-66. (IHC)
- 4. del Rio ML et al. (2011) Flt3L-mobilized dendritic cells bearing H2-Kbm1 apoptotic cells do not induce cross-tolerance to CD8+ T cells across a class I MHC mismatched barrier. Transpl Int 24(5): 501–13. (FC)
- 5. Mitchell LM et al. (2011) Distinct responses of splenic dendritic cell subsets to infection with Listeria monocytogenes: maturation phenotype, level of infection, and T cell priming capacity ex vivo. Cell Immunol 268(2): 79–86. (FC)
- 6. Charles N et al. (2010) Basophils and the T helper 2 environment can promote the development of lupus nephritis. Nat Med. 16(6): 701–07. (FC)
- 7. McKee AS et al. (2009) Alum induces innate immune responses through macrophage and mast cell sensors, but these sensors are not required for alum to act as an adjuvant for specific immunity. J Immunol 183(7): 4403–14.
- 8. Schuhmann B et al. (2005) A role for brain-derived neurotrophic factor in B cell development. J Neuroimmunol 163(1-2): 15–23. (IHC)
- 9. Rosmalen JG et al. (2000) Subsets of macrophages and dendritic cells in nonobese diabetic mouse pancreatic inflammatory infiltrates: Correlation with the development of diabetes. Lab Invest 80(1): 23–30.
- 10. Monteith CE et al. (1996) Identification of monoclonal antibodies for immunohistochemical staining of feline B lymphocytes in frozen and formalin-fixed paraffin-embedded tissues. Can J Vet Res 60(3): 193–8. (IHC)
- 11. Rolink A et al. (1996) A subpopulation of B220+ cells in murine bone marrow does not express CD19 and contains natural killer cell progenitors. J Exp Med 183(1): 187–94.
- 12. Whiteland JL et al. (1995) Immunohistochemical detection of T Cell subsets and other leucocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem 43(3): 313–20. (IHC)
- 13. Watanabe Y & Akaike T. (1994) Activation signal induces the expression of B cell-specific CD45R epitope (6B2) on murine T cells. Scand J Immunol. 39(5): 419–25.
- 14. Johnson P et al. (1989) Identification of the alternatively spliced exons of murine CD45 (T200) required for reactivity with B220 and other T200-restricted antibodies. J Exp Med 169(3): 1179–84.
- 15. Coffman RL. (1982) Surface antigen expression and immunoglobulin gene rearrangement during mouse pre- B cell development. Immunol Rev 69: 5–23. (IP)
- 16. Morse H et al. (1982) Abnormalities induced by the mutant gene lpr; Expansion of a unique lymphocyte subset. J Immunol 129(6): 2612-5.
- 17. Coffman RL & Weissman IL. (1981) B220: A B cell-specific member of the T200 glycoprotein family. Nature 289(5799): 681-3.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. Alexa Fluor and Pacific Blue are trademarks of Life Technologies Corporation. Antibodies conjugated to Alexa Fluor® or Pacific Blue™ are licensed for internal research use only and sale is expressly conditioned on the buyer not using the antibody for manufacturing, performing a service or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.