

Small Molecules

HA-100

Protein kinase inhibitor; Inhibits PKA, PKC, and PKG

Catalog # 72482
72484

5 mg
25 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

HA-100 is an isoquinoline compound with an added piperazinylsulfonyl group that acts as an inhibitor of protein kinases (PKs), including PKA, PKC, and PKG (IC_{50} = 8, 12, and 4 μ M, respectively; Hagiwara et al.). It less effectively blocks the activity of myosin light chain kinase (IC_{50} = 240 μ M; Hagiwara et al.). This product is supplied as the dihydrochloride salt of the molecule.

Molecular Name:	HA-100 (Dihydrochloride)
Alternative Names:	Not applicable
CAS Number:	210297-47-5
Chemical Formula:	$C_{13}H_{15}N_3O_2S \cdot 2HCl$
Molecular Weight:	350.3 g/mol
Purity:	$\geq 95\%$
Chemical Name:	C-1; 5-(1-piperazinylsulfonyl)-isoquinoline, dihydrochloride
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	\cdot DMSO ≤ 3.1 mM For example, to prepare a 2 mM stock solution in DMSO, resuspend 1 mg in 1.43 mL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

MAINTENANCE AND SELF-RENEWAL

- Improves single cell survival and supports high cloning efficiency in human pluripotent stem cell cultures (Chen et al.).

REPROGRAMMING

- Increases human fibroblast reprogramming efficiency with PD0325901, CHIR99021, A83-01 and hLIF (Yu et al.).

References

Chen G et al. (2011) Chemically defined conditions for human iPSC derivation and culture. Nat Methods 8(5): 424–9.

Hagiwara M et al. (1987) Selective modulation of calcium-dependent myosin phosphorylation by novel protein kinase inhibitors, isoquinolinesulfonamide derivatives. Mol Pharmacol 32(1): 7–12.

Yu J et al. (2011) Efficient feeder-free episomal reprogramming with small molecules. M. Pera (Ed.). PLoS One 6(3): e17557.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.