OAC1

Small Molecules

Inducer of OCT4 expression

STEMCELL'

Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 72602 5 mg

Product Description

OAC1 is an OCT4-activating compound that activates expression through the OCT4 gene promoter (Li et al.). OCT4 (POU5F1) is a transcription factor that is critically involved in the self-renewal of pluripotent stem cells, and its expression is commonly used as a marker for pluripotency. With SOX2, KLF4, and c-MYC, OCT4 is involved in the reprogramming of somatic cells to produce induced pluripotent stem cells (Niwa et al.; Takahashi et al.).

Molecular Name: OAC1

Alternative Names: Not applicable CAS Number: 300586-90-7 Chemical Formula: $C_{14}H_{11}N_3O$ Molecular Weight: 237.3 g/mol Purity: $\geq 98\%$

Chemical Name: N-1H-pyrrolo[2,3-c]pyridin-5-yl-benzamide

Structure:

Properties

Physical Appearance: A crystalline solid

Storage: Product stable at -20°C as supplied. Protect from prolonged exposure to light.

Stable as supplied for 12 months from date of receipt.

Solubility: · Absolute ethanol ≤ 10 mM

· DMSO ≤ 42 mM

For example, to prepare a 10 mM stock solution in DMSO, resuspend 5 mg in 2.11 mL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Small Molecules OAC1



Published Applications

MAINTENANCE AND SELF-RENEWAL

- · Mediates ex vivo expansion of cord blood CD34+ hematopoietic stem and progenitor cells (Huang et al.). REPROGRAMMING
- · Enhances the reprogramming efficiency of mouse embryonic fibroblasts transfected with OCT4, SOX2, KLF4, and c-MYC (Li et al.).

References

Huang X et al. (2016) Activation of OCT4 enhances ex vivo expansion of human cord blood hematopoietic stem and progenitor cells by regulating HOXB4 expression. Leukemia 30(1): 144–53.

Li W et al. (2012) Identification of Oct4-activating compounds that enhance reprogramming efficiency. Proc Natl Acad Sci USA 109(51): 20853–8.

Niw H et al. (2000) Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. Nat Genet 24(4): 372–6.

Takahasi K et al. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell 131(5): 861–72.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.