Gatifloxacin

Small Molecules

Antibiotic; Inhibits bacterial DNA gyrase and topoisomerase IV

Catalog # 72752 1 g



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Gatifloxacin is a fluoroquinolone antibiotic which inhibits bacterial DNA gyrase ($IC_{50} = 0.109 \,\mu\text{g/ml}$) and topoisomerase IV ($IC_{50} = 13.8 \,\mu\text{g/ml}$; Takei et al.). It is much less effective against HeLa cell topoisomerase II ($IC_{50} = 265 \,\mu\text{g/ml}$; Takei et al.).

Molecular Name: Gatifloxacin

Alternative Names: AM 1155; BMS 206584-01; PD 135432

CAS Number: 112811-59-3 Chemical Formula: $C_{19}H_{22}FN_3O_4$ Molecular Weight: 375.4 g/mol Purity: \geq 98%

Chemical Name: 1-Cyclopropyl-6-fluoro-8-methoxy-7-(3-methylpiperazin-1-yl)-4-oxo-1,4-dihydroquinoline-3-carboxylic acid

Structure:

Properties

Physical Appearance: A crystalline solid

Storage: Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please

contact techsupport@stemcell.com.

Solubility: · Absolute ethanol ≤ 20 mM

· DMSO ≤ 45 mM

For example, to prepare a 20 mM stock solution in DMSO, resuspend 1 mg in 225 µL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Small Molecules Gatifloxacin



Published Applications

MAINTENANCE AND SELF-RENEWAL

· Promotes self-renewal in cultured human and mouse embryonic stem (ES) cells (Desbordes et al.).

References

Desbordes SC et al. (2008) High-throughput screening assay for the identification of compounds regulating self-renewal and differentiation in human embryonic stem cells. Cell Stem Cell 2(6): 602–12.

Takei M et al. (1998) Inhibitory Activities of Gatifloxacin (AM-1155), a Newly Developed Fluoroquinolone, against Bacterial and Mammalian Type II Topoisomerases. Antimicrob Agents Chemother 42(10): 2678–2681.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.