

Small Molecules

IBMX

cAMP pathway activator; Inhibits cyclic nucleotide phosphodiesterases

Catalog # 72762
72764

100 mg
500 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

IBMX is a widely-used non-specific inhibitor of cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) phosphodiesterases (PDEs; IC_{50} = 19, 50, 18, 13, 32, 7, and 50 μ M for PDE1, PDE2, PDE3, PDE4, PDE5, PDE7, and PDE11, respectively). PDE8A, PDE8B, and PDE9 are insensitive to IBMX (Fawcett et al.). By inhibiting PDEs, IBMX increases cellular cAMP and cGMP levels, activating cyclic-nucleotide-regulated protein kinases. Methylxanthines, including IBMX, caffeine, and theophylline, bind adenosine receptors, typically antagonizing the suppressive effects of natural agonists (Snyder et al.).

Molecular Name:	IBMX
Alternative Names:	1-Methyl-3-Isobutylxanthine; Isobutylmethylxanthin; Isobutyl methylxanthine; NSC 165960
CAS Number:	28822-58-4
Chemical Formula:	$C_{10}H_{14}N_4O_2$
Molecular Weight:	222.3 g/mol
Purity:	$\geq 98\%$
Chemical Name:	3,7-dihydro-3-isobutyl-1-methyl-1H-purine-2,6-dione
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	<ul style="list-style-type: none">· Absolute ethanol ≤ 20 mM· DMSO ≤ 40 mM For example, to prepare a 10 mM stock solution in DMSO, resuspend 10 mg in 4.5 mL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C . Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Used in combination with fibroblast growth factor (FGF) 1, dopamine, 12-O-tetradecanoylphorbol-13-acetate (TPA), and forskolin to induce expression of the dopaminergic neuron marker tyrosine hydroxylase in neurons derived from the human NT2 cell line (Iacovitti et al.).
- Used in combination with dexamethasone, insulin, and indomethacin for in vitro induction of adipogenic differentiation of unrestricted somatic stem cells (USSCs), a CD45-negative population of stem cells isolated from human cord blood (Kögler et al.; Pittenger et al.).
- Induces neural differentiation from human umbilical cord blood-derived mesenchymal stem cells (MSCs; Tio et al.).
- Promotes the differentiation of rat neural progenitor cells (NPCs) into functional neurons in vitro (Lepski et al.).

References

- Fawcett L et al. (2000) Molecular cloning and characterization of a distinct human phosphodiesterase gene family: PDE11A. *Proc Natl Acad Sci* 97(7): 3702–3707.
- Iacovitti L et al. (2001) Differentiation of human dopamine neurons from an embryonic carcinomal stem cell line. *Brain Res* 912(1): 99–104.
- Kögler G et al. (2004) A new human somatic stem cell from placental cord blood with intrinsic pluripotent differentiation potential. *J Exp Med* 200(2): 123–35.
- Lepski G et al. (2013) cAMP promotes the differentiation of neural progenitor cells in vitro via modulation of voltage-gated calcium channels. *Front Cell Neurosci* 7: 155.
- Pittenger MF et al. (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284(5411): 143–7.
- Snyder SH et al. (1981) Adenosine receptors and behavioral actions of methylxanthines. *Proc Natl Acad Sci U S A* 78(5): 3260–4.
- Tio M et al. (2010) Roles of db-cAMP, IBMX and RA in aspects of neural differentiation of cord blood derived mesenchymal-like stem cells. *PLoS One* 5(2): e9398.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.