

Small Molecules

Pifithrin-mu

p53 inhibitor

Catalog # 72802
72804

10 mg
50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Pifithrin-mu (PFT- μ) is an inhibitor of p53-mediated apoptosis, preventing p53 binding to Bcl-xL and Bcl-2 at the mitochondrial surface, without affecting p53 transactivational activities (Strom et al.). In vitro, PFT- μ binds both p53 (K_d = 0.82 mM) and Bcl-xL (K_d = 0.80 mM; Hagn et al.). PFT- μ also interacts selectively with heat shock protein 70 (HSP70), leading to disruption of the association between HSP70 and many of its co-chaperones and substrate proteins (Leu et al.).

Molecular Name:	Pifithrin-mu
Alternative Names:	2-Phenylethynylsulfonamide; PFT- μ ; Pifithrin- μ
CAS Number:	64984-31-2
Chemical Formula:	$C_8H_7NO_2S$
Molecular Weight:	181.2 g/mol
Purity:	$\geq 98\%$
Chemical Name:	Ethynylsulfonamide, 2-phenyl-
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	· Absolute ethanol ≤ 75 mM · DMSO ≤ 75 mM For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 552 μ L of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

MAINTENANCE AND SELF-RENEWAL

- In combination with Rho-associated coiled-coil containing protein kinase (ROCK) inhibitor Y-27632, improves cell recovery after cryopreservation (Xu et al.).
- Inhibits DNA damage-induced apoptosis in human embryonic stem (ES) cells (Qin et al.).

References

- Hahn F et al. (2010) BclxL changes conformation upon binding to wild-type but not mutant p53 DNA binding domain. J Biol Chem 285(5): 3439–50.
- Leu JI-J et al. (2009) A small molecule inhibitor of inducible heat shock protein 70. Mol Cell 36(1): 15–27.
- Qin H et al. (2007) Regulation of apoptosis and differentiation by p53 in human embryonic stem cells. J Biol Chem 282(8): 5842–52.
- Strom E et al. (2006) Small-molecule inhibitor of p53 binding to mitochondria protects mice from gamma radiation. Nat Chem Biol 2(9): 474–9.
- Xu X et al. (2010) Enhancement of cell recovery for dissociated human embryonic stem cells after cryopreservation. Biotechnol Prog 26(3): 781–8.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.