

Small Molecules

PluriSIn-1

Oleic acid biosynthesis pathway inhibitor; Inhibits stearyl-CoA desaturase (SCD1)

Catalog # 72822
72824

10 mg
50 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

PluriSIn-1 is an N-acyl phenylhydrazine derivative that inhibits stearyl-CoA desaturase, a key enzyme for lipid metabolism that is expressed in human pluripotent stem cells (Ben-David et al.).

Molecular Name:	PluriSIn-1
Alternative Names:	N'-phenyl-hydrazine-Isonicotinic acid; NSC 14613
CAS Number:	91396-88-2
Chemical Formula:	C ₁₂ H ₁₁ N ₃ O
Molecular Weight:	213.2 g/mol
Purity:	≥ 95%
Chemical Name:	N'-phenylpyridine-4-carbohydrazide
Structure:	



Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Solubility:	<ul style="list-style-type: none">· Absolute ethanol ≤ 90 mM· DMSO ≤ 140 mM For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 469 µL of fresh DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Selectively eliminates undifferentiated human embryonic stem (ES) and induced pluripotent stem (iPS) cells while sparing differentiated cells, and prevents teratoma formation in transplanted mice (Ben-David et al.).
- Induces apoptosis of Nanog-positive iPS cells in vitro, while leaving iPS cell-derived cardiomyocytes unaffected (Zhang et al.).

References

Ben-David U et al. (2013) Selective elimination of human pluripotent stem cells by an oleate synthesis inhibitor discovered in a high-throughput screen. *Cell Stem Cell* 12(2): 167–79.

Zhang L et al. (2014) Inhibition of stearyl-coA desaturase selectively eliminates tumorigenic Nanog-positive cells: improving the safety of iPS cell transplantation to myocardium. *Cell Cycle* 13(5): 762–71.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.