#### U-0126

# **Small Molecules**

MEK/ERK pathway inhibitor; Inhibits

MEK1 and MEK2

Catalog # 73522 1 mg 73524

10 mg



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## **Product Description**

U-0126 is a selective, non-ATP competitive inhibitor of mitogen activated protein kinase kinase (MEK), inhibiting MEK1 and MEK2 with IC<sub>50</sub> values of 72 nM and 58 nM, respectively (Favata et al.; Scherle et al.). It shows little or no inhibition at micromolar levels of other kinases such as ERK, protein kinase C (PKC), c-Jun N-terminal kinases (JNK), other MAP kinase kinases (MKK3, MKK4, MKK6), cyclindependant kinases (CDK2, CDK4), ABL, and RAF (Favata et al. 1998). U-0126 also antagonizes AP-1 transcription and selectively inhibits promoters containing AP-1 response elements (Favata et al.).

Molecular Name: U-0126

Alternative Names: Not applicable CAS Number: 109511-58-2 Chemical Formula:  $C_{18}H_{16}N_6S_2$ Molecular Weight: 380.5 g/mol Purity: ≥ 98%

Chemical Name: 2,3-bis[amino[(2-aminophenyl)thio]methylene]-butanedinitrile

Structure:

### **Properties**

Physical Appearance:

A crystalline solid

Storage:

Product stable at -20°C as supplied. Protect from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com.

Solubility:

· DMSO ≤ 65 mM

· Absolute ethanol ≤ 1 mM

For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 263 µL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

## Small Molecules U-0126



### **Published Applications**

MAINTENANCE AND SELF-RENEWAL

- · In combination with FGF2, Activin A, and a PKC inhibitor, U-0126 promotes maintenance of human pluripotent stem cells (Kinehara et al.).
- · Used alone with MEF-conditioned medium, U-0126 inhibits self-renewal of human pluripotent stem cells, leading to differentiation, without affecting proliferation or survival (Li et al.).
- · Inhibits glutamate-induced oxidative stress in mouse hippocampal HT22 cells and rat primary cortical cultures (Satoh et al.; Ong et al.).
- · Neuroprotective in a gerbil ischemia model and in primary mouse neurons cultured under hypoxia (Namura et al.). IMMUNOLOGY
- · Inhibits T-cell proliferation against certain antigenic stimuli by reducing IL-2 mRNA levels (DeSilva et al.). DISEASE MODELING
- · Activates peroxisome proliferator activated receptor (PPAR) co-activator 1 $\alpha$  (PGC-1 $\alpha$ ) and prevents neurotoxicity and spatial memory impairment in rats challenged by amyloid beta (A $\beta$ ; Ashabi et al.).

#### References

Ashabi G et al. (2012) ERK and p38 inhibitors attenuate memory deficits and increase CREB phosphorylation and PGC-1 $\alpha$  levels in A $\beta$ -injected rats. Behav Brain Res 232(1): 165–73.

DeSilva DR et al. (1998) Inhibition of mitogen-activated protein kinase kinase blocks T cell proliferation but does not induce or prevent anergy. J Immunol 160(9): 4175–81.

Favata MF et al. (1998) Identification of a novel inhibitor of mitogen-activated protein kinase kinase. J Biol Chem 273(29): 18623–32. Kinehara M et al. (2013) Protein kinase C regulates human pluripotent stem cell self-renewal. PLoS One 8(1): e54122.

Li J et al. (2007) MEK/ERK signaling contributes to the maintenance of human embryonic stem cell self-renewal. Differentiation 75(4): 299–307.

Namura S et al. (2001) Intravenous administration of MEK inhibitor U0126 affords brain protection against forebrain ischemia and focal cerebral ischemia. Proc Natl Acad Sci U S A 98(20): 11569–74.

Ong Q et al. (2015) U0126 protects cells against oxidative stress independent of its function as a MEK inhibitor. ACS Chem Neurosci 6(1): 130–7.

Satoh T et al. (2000) Neuroprotection by MAPK/ERK kinase inhibition with U0126 against oxidative stress in a mouse neuronal cell line and rat primary cultured cortical neurons. Neurosci Lett 288(2): 163–6.

Scherle PA et al. (1998) Inhibition of MAP kinase kinase prevents cytokine and prostaglandin E2 production in lipopolysaccharide-stimulated monocytes. J Immunol 161(10): 5681–6.

#### Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.