# Anti-Mouse CD45.1 Antibody, Clone A20, Biotin

# **Antibodies**

Mouse monoclonal IgG2a antibody against mouse CD45.1, biotin-

conjugated

Catalog #60117BT #60117BT.1

500 μg 0.5 mg/mL

50 μg 0.5 mg/mL



Scientists Helping Scientists<sup>™</sup> | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

# **Product Description**

The A20 monoclonal antibody reacts with mouse alloantigen CD45.1 (Ly5.1) expressed by all hematopoietic cells except mature erythrocytes and platelets in mouse strains RIII, SJL/J, STS/A, and DA. Clone A20 does not react with leukocytes that express CD45.2 alloantigen. CD45.1 is an allelic form of CD45 (Leukocyte Common Antigen), a 180 - 240 kD type I transmembrane glycoprotein. CD45 is a member of the protein tyrosine phosphatase (PTP) family and contains two catalytic domains in the intracellular (COOH) domain, which regulate cellular signaling. The extracellular domains are highly variable, due to alternative splicing among exons 4, 5, and 6, and differential glycosylation. These CD45 isoforms correlate to specific cell types, as well as specific activation and maturation states of immune cells. The A20 monoclonal antibody has been used extensively in adoptive cell transfer and bone marrow transplantation studies in mice, allowing for the differentiation of CD45.2- and CD45.1- expressing cells.

Target Antigen Name: CD45.1

Alternative Names: B220, CD45 antigen, CD45R, GP180, Leukocyte common antigen (LCA), Ly5.1, Lyt4, Protein tyrosine

phosphatase receptor type C (Ptprc), RT7.1, T200

Gene ID: 19264

Species Reactivity: Mouse, does not react with mouse CD45.2

Host Species: Mouse (A.SW)
Clonality: Monoclonal

Clone: A20

Isotype: IgG2a, kappa

Immunogen: SJL mouse thymocytes and splenocytes

Conjugate: Biotin

# **Applications**

Verified: FC

Reported: FC, IHC, IF

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

# **Properties**

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide

Purification: The antibody was purified by affinity chromatography and conjugated with biotin under optimal conditions.

The solution is free of unconjugated biotin.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact

techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is ≤ 0.25 µg per 1 x 10^6 cells in 100 µL volume. It is


recommended that the antibody be titrated for optimal performance for each application.

#### Anti-Mouse CD45.1 Antibody, Clone A20, Biotin

# **Antibodies**



### Data



Flow cytometry analysis of SLJ mouse splenocytes labeled with Anti-Mouse CD45.1 Antibody, Clone A20, Biotin followed by streptavidin (SAV) PE (filled histogram).

### Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

### References

- 1. Rabenhorst U et al. (2015) Single-stranded DNA-binding transcriptional regulator FUBP1 is essential for fetal and adult hematopoietic stem cell self-renewal. Cell Rep 11(12): 1847–55. (FACS, FC)
- 2. Van Blijswijk J et al. (2015) Altered lymph node composition in diphtheria toxin receptor-based mouse models to ablate dendritic cells. J Immunol 194(1): 307–15. (FC)
- 3. Gebhardt T et al. (2009) Memory T cells in nonlymphoid tissue that provide enhanced local immunity during infection with herpes simplex virus. Nat Immunol 10(5): 524–30. (FC)
- 4. Trowbridge JJ et al. (2009) DNA methyltransferase 1 is essential for and uniquely regulates hematopoietic stem and progenitor cells. Cell Stem Cell 5(4): 442–9. (FC)
- 5. Mende I et al. (2006) Flk2+ myeloid progenitors are the main source of Langerhans cells. Blood 107(4): 1383-90. (FC, IF)
- 6. Duncan AW et al. (2005) Integration of Notch and Wnt signaling in hematopoietic stem cell maintenance. Nat Immunol 6(3): 314-22. (FC)
- 7. Hock H et al. (2004) Gfi-1 restricts proliferation and preserves functional integrity of haematopoietic stem cells. Nature 431(7011): 1002-7. (FC)
- 8. Lessner SM et al. (2002) Atherosclerotic lesions grow through recruitment and proliferation of circulating monocytes in a murine model. Am J Pathol 160(6): 2145–55. (FC, IF, IHC)
- 9. Werner N et al. (2002) Bone marrow-derived progenitor cells modulate vascular reendothelialization and neointimal formation: effect of 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibition. Arterioscler Thromb Vasc Biol 22(10): 1567–72. (IHC)
- 10. Wilson A et al. (2001) Notch 1-deficient common lymphoid precursors adopt a B cell fate in the thymus. J Exp Med 194(7): 1003–12. (FC)
- 11. Shen FW et al. (1986) Further definition of the Ly-5 system. Immunogenetics 24(3): 146-9.
- 12. Yakura H et al. (1983) On the function of Ly-5 in the regulation of antigen-driven B cell differentiation. Comparison and contrast with Lyb-2. J Exp Med 157(4): 1077–88.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, Scientists Helping Scientists, and EasySep are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. Alexa Fluor® is a registered trademark of Life Technologies Corporation. This product is licensed for internal research use only and its sale is expressly conditioned on the buyer not using it for manufacturing, performing a service, or medical test, or otherwise generating revenue. For use other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.