

Cytokines

Mouse Recombinant GM-CSF

Granulocyte-macrophage colony-stimulating factor



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	78017.1	20 µg
	78017	100 µg
	78017.2	1000 µg

Product Description

Granulocyte-macrophage colony-stimulating factor (GM-CSF) promotes the proliferation and differentiation of hematopoietic progenitor cells and the generation of neutrophils, eosinophils, and macrophages. In synergy with other cytokines such as stem cell factor, IL-3, erythropoietin, and thrombopoietin, it also stimulates erythroid and megakaryocyte progenitors (Barreda et al.). GM-CSF was first purified from the culture of mouse lung tissue after lipopolysaccharide treatment. GM-CSF is produced by multiple cell types, including stromal cells, Paneth cells, macrophages, dendritic cells (DCs), endothelial cells, smooth muscle cells, fibroblasts, chondrocytes, and Th1 and Th17 cells (Francisco-Cruz et al.). The receptor for GM-CSF (GM-CSFR) is composed of two subunits: the cytokine-specific α subunit (GMR α ; CD116) and the common subunit β c (CD131) shared with IL-3 and IL-5 receptors (Broughton et al.). GM-CSFR is expressed on hematopoietic cells, including progenitor cells and immune cells, as well as non-hematopoietic cells. GM-CSF is able to stimulate the development of DCs that ingest, process, and present antigens to the immune system (Francisco-Cruz et al.).

Product Information

Alternative Names:	Colony-stimulating factor 2, CSF-2, MGI-1GM, Pluripoietic-alpha
Accession Number:	P01587
Amino Acid Sequence:	MAPTRSPITV TRPWKHVEAI KEALNLLDDM PVTLNEEVEV VSNEFSFKKL TCVQTRLKIF EQGLRGNFTK LKGALNMTAS YYQTYCPPTP ETD CETQVTT YADFIDSLKT FLTDIPFECK KPVQK
Predicted Molecular Mass:	14.3 kDa
Species:	Mouse
Cross Reactivity:	Highly species-specific
Formulation:	Lyophilized from a sterile filtered aqueous solution containing acetic acid.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 2 \times 10^7$ units/mg ($EC_{50} \leq 0.05$ ng/mL) as determined by a cell proliferation assay using FDC-P1 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage:	Store at -20°C to -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Reconstitution:	Centrifuge vial before opening. Resuspend the product in sterile water containing 0.1% bovine serum albumin (BSA) to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 month or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles. NOTE: If reconstituted product will be used immediately BSA is not required.

Data



(A) The biological activity of Mouse Recombinant GM-CSF was tested by its ability to promote the proliferation of FDC-P1 cells. Cell proliferation was measured after 91 hours of culture using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC50 in the above example is 3 - 5 pg/mL.

(B) 1 μ g of Mouse Recombinant GM-CSF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant GM-CSF has a predicted molecular mass of 14.3 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Barreda DR et al. (2004) Regulation of myeloid development and function by colony stimulating factors. *Dev Comp Immunol* 28(5): 509–54.
- Broughton SE et al. (2012) The GM-CSF/IL-3/IL-5 cytokine receptor family: from ligand recognition to initiation of signaling. *Immunol Rev* 250(1): 277–302.
- Francisco-Cruz A et al. (2014) Granulocyte-macrophage colony-stimulating factor: not just another haematopoietic growth factor. *Med Oncol* 31(1): 774.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.