

Cytokines

Human Recombinant IFN-gamma

Interferon-gamma



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	78020.1	50 µg
	78020	100 µg
	78020.2	1000 µg

Product Description

Interferon-gamma (IFN- γ), also known as type II interferon, is produced by T and NK cells, and in smaller amounts by dendritic cells and macrophages. IFN- γ is controlled by cytokines such as IL-12 and IL-18 secreted in response to infection (Schroder et al.). IFN- γ binds to a receptor complex and initiates signal transduction via the JAK/STAT pathway; this culminates in the transcription and activation of many genes that control a diverse array of immunological functions (de Weerd & Nguyen; Krause et al.). IFN- γ stimulates the antimicrobial and anti-tumor activity of macrophages, NK cells, and neutrophils (Billiau & Matthys) by promoting the activation of microbial effector functions such as production of reactive oxygen species, nitric oxide, and complement (Schroder et al.). IFN- γ enhances MHC class I and II expression in dendritic cells and mononuclear phagocytes, as well as the production of IL-12 by dendritic cells. In B cells, IFN- γ stimulates survival and growth in both mouse and human cells, and redirects B cells from proliferation towards differentiation. IFN- γ favors the development of Th1 vs Th2 cells and stimulates monocyte differentiation and function (Schroder et al.).

Product Information

Alternative Names:	Interferon gamma, Type II interferon
Accession Number:	P01579
Amino Acid Sequence:	MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI VSFYFKLFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS VTDLNVQRKA IHELIQVMAE LSPAAGTKGR KRSQMLFRGR RASQ
Predicted Molecular Mass:	17 kDa
Species:	Human
Cross Reactivity:	Mouse, Monkey
Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 2 \times 10^7$ units/mg ($EC_{50} \leq 0.05$ ng/mL) as determined by a cytotoxicity assay using HT-29 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 0.2 EU/ μ g protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water or phosphate-buffered saline to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 week or at -20°C to -80°C for up to 2 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant IFN-gamma was tested by its ability to promote the cytotoxicity of HT-29 (HTB-38) cells. Cytotoxicity was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell death is at 50% of maximum. The EC₅₀ for this effect is 0.035 ng/mL.

(B) 2 µg of Human Recombinant IFN-gamma was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant IFN-gamma has a predicted molecular mass of 17 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Billiau A & Matthys P. (2009) Interferon-gamma: a historical perspective. *Cytokine Growth Factor Rev* 20(2): 97–113.
- de Weerd NA & Nguyen T. (2012) The interferons and their receptors--distribution and regulation. *Immunol Cell Biol* 90(5): 483–91.
- Krause CD et al. (2000) Signaling by covalent heterodimers of interferon-gamma. Evidence for one-sided signaling in the active tetrameric receptor complex. *J Biol Chem* 275(30): 22995–3004.
- Schroder K et al. (2004) Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 75(2): 163–89.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.