Cytokines Mouse Recombinant IFN-gamma

Interferon-gamma



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLORAL CONTACT DETAILS VISIT OUR WERSITE

Catalog # 78021.1 20 μg

78021 100 μg 78021.2 1000 μg

Product Description

Interferon-gamma (IFN- γ), also known as type II interferon, is produced by T and NK cells, and in smaller amounts by dendritic cells and macrophages. IFN- γ is controlled by cytokines such as IL-12 and IL-18 secreted in response to infection (Schroder et al.). IFN- γ binds to a receptor complex and initiates signal transduction via the JAK/STAT pathway; this culminates in the transcription and activation of many genes that control a diverse array of immunological functions (De Weerd and Nguyen; Krause et al.). IFN- γ stimulates the antimicrobial and anti-tumor activity of macrophages, NK cells, and neutrophils (Billiau & Matthys) by promoting the activation of microbial effector functions such as production of reactive oxygen species, NO intermediates, complement, etc. (Schroder et al.). IFN- γ enhances MHC class I and II expression in dendritic cells and mononuclear phagocytes, as well as the production of IL-12 by dendritic cells. In B cells, IFN- γ stimulates survival and growth in both mouse and human cells, and redirects B cells from proliferation towards differentiation. IFN- γ favors the development of Th1 vs Th2 cells and stimulates monocyte differentiation and function (Schroder et al.).

Product Information

Alternative Names: Interferon gamma, Type II interferon

Accession Number: P01580

Amino Acid Sequence: MHGTVIESLE SLNNYFNSSG IDVEEKSLFL DIWRNWQKDG DMKILQSQII SFYLRLFEVL KDNQAISNNI

SVIESHLITT FFSNSKAKKD AFMSIAKFEV NNPQVQRQAF NELIRVVHQL LPESSLRKRK RSRC

Predicted Molecular Mass: 15.7 kDa

Species: Mouse

Cross Reactivity: Human, Rat

Formulation: Lyophilized from a sterile filtered aqueous solution containing sodium phosphate and sodium chloride, pH

7.5.

Source: E. coli

Specifications

Activity: The specific activity is ≥ 1 x 10^6 units/mg as determined in a viral challenge assay using EMC virus on

L929 cells.

Purity: $\geq 95\%$

Endotoxin Level: Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage: Store at -20°C to -80°C.

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Resuspend the product in sterile water containing 0.1% bovine serum

albumin (BSA) to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 month or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.


NOTE: If reconstituted product will be used immediately BSA is not required.

Cytokines

Mouse Recombinant IFN-gamma



Data



(A) The biological activity of Mouse Recombinant IFN-gamma was measured with L929 cells exposed to the encephalomyocarditis (EMC) virus in a cytopathic effect (CPE) assay. The specific activity in the above example is 1.25 x 10^6 units/mg.

(B) 1 µg of Mouse Recombinant IFN-gamma was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant IFN-gamma has a predicted molecular mass of 15.7 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Billiau A & Matthys P. (2009) Interferon-gamma: a historical perspective. Cytokine Growth Factor Rev 20(2): 97–113. Krause CD et al. (2000) Signaling by covalent heterodimers of interferon-gamma. Evidence for one-sided signaling in the active tetrameric receptor complex. J Biol Chem 275(30): 22995–3004.

Schroder K et al. (2004) Interferon-gamma: an overview of signals, mechanisms and functions. J Leukoc Biol 75(2): 163–89. de Weerd NA & Nguyen T. (2012) The interferons and their receptors--distribution and regulation. Immunol Cell Biol 90(5): 483–91.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.