

Cytokines

Human Recombinant Noggin



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Noggin

Catalog #	78060.1	10 µg
	78060	50 µg
	78060.2	1000 µg

Product Description

Noggin binds to and antagonizes bone morphogenic protein (BMP) ligands of the transforming growth factor - beta (TGF-β) family. Noggin supports maintenance of undifferentiated human embryonic stem cells in vitro, and can be used to prevent spontaneous differentiation in the short-term (Chaturvedi et al.). Noggin is essential for development of structures derived from ectoderm embryonic somite, skeletal patterning, and neurogenesis in vivo. It also influences chondrogenesis, osteogenesis and joint formation (Krause et al.), and promotes dopaminergic differentiation of embryonic stem cells and subsequent survival of dopamine neurons (Chiba et al.).

Product Information

Alternative Names:	NOG, SYM1, SYNS1, Symphalangism 1 (proximal), Synostoses (multiple) syndrome
Accession Number:	Q13253
Amino Acid Sequence:	QHYLEHIRPAP SDNLPLVDLI EHPDPIFDPK EKDLNETLLR SLLGGHYDPG FMATSPPEDR PGGGGGAAGG AEDLAELDQL LRQRPSGAMP SEIKGLEFSE GLAQGKKQRL SKKLRRKLQM WLWSQTFCPV LYAWNDLGSR FWPRYVKVGS CFSKRSCSVP EGMVCKPSKS VHLTTLRWRC QRRGGQRCGW IPIQYPIISE CKCSC
Predicted Molecular Mass:	30 kDa
Species:	Human
Cross Reactivity:	Mouse
Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	CHO


Specifications

Activity:	The specific activity is $> 4 \times 10^5$ units/mg ($EC_{50} < 2.5$ ng/mL) as determined by a bioassay using ATDC5 cells in the presence of 10 ng/mL of human BMP-4.
Purity:	$> 95\%$
Endotoxin Level:	Measured by kinetic limulus amebocyte lysate (LAL) analysis and is < 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C .
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Resuspend the product in sterile water or phosphate-buffered saline to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at $2 - 8^{\circ}\text{C}$ for up to 1 week or at -20°C to -80°C for up to 2 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant Noggin was tested by its ability to inhibit BMP-4 induced alkaline phosphatase production of ATDC-5 cells. Inhibition of BMP-4 induced alkaline phosphatase production was measured using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which inhibition of alkaline phosphatase production is at 50% of maximum. The EC50 in the above example is 1.9 ng/mL.

(B) 2 µg of Human Recombinant Noggin was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant Noggin has a predicted molecular mass of 30 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Chaturvedi G et al. (2009) Noggin maintains pluripotency of human embryonic stem cells grown on Matrigel. *Cell Prolif* 42(4): 425–33.
- Chiba S et al. (2008) Noggin enhances dopamine neuron production from human embryonic stem cells and improves behavioral outcome after transplantation into Parkinsonian rats. *Stem Cells* 26(11): 2810–20.
- Krause C et al. (2011) Noggin. *Int J Biochem Cell Biol* 43(4): 478–81.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.