# Anti-Human CD45 Antibody, Clone 2D1, FITC

### **Antibodies**

Mouse monoclonal IgG1 antibody against human CD45, FITC-conjugated

Catalog #60123FI 100 Tests 5 μL/test #60123FI.1 25 Tests 5 μL/test



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## **Product Description**

The 2D1 antibody reacts with all isoforms of CD45, a type I transmembrane glycoprotein expressed on the surface of hematopoietic stem and progenitor cells and mature blood cells, except mature erythrocytes, platelets, and plasma cells; expression of CD45 is lost during differentiation of these cell lineages. CD45 is a member of the protein tyrosine phosphatase family and functions in a number of immunoregulatory processes, including cell activation, growth, differentiation, and oncogenic transformation. The large cytoplasmic portion of CD45 contains two tyrosine phosphatase domains, one of which is enzymatically active, that are involved in modulating the function of intracellular substrates such as the Src kinases Lck and Fyn. Several isoforms of CD45 have been identified, the expression of which differs according to cell type and functional status. Alternative splicing of three exons (4, 5, and 6) encoding the extracellular RA, RB, and RC polypeptide sequences gives rise to up to 8 isoforms with molecular masses in the range of 180 - 240 kDa. The leukocyte common antigen, the region recognized by the 2D1 antibody, is an extracellular region proximal to the membrane and common to all isoforms of CD45.

Target Antigen Name: CD45

Alternative Names: Hle-1, LCA, Leukocyte common antigen, Ly-5, T200

Gene ID: 5788
Species Reactivity: Human

Host Species: Mouse (BALB/c)
Clonality: Monoclonal

Clone: 2D1

Isotype: IgG1, kappa

Immunogen: Human peripheral blood mononuclear cells

Conjugate: FITC (Fluorescein isothiocyanate)

# **Applications**

Verified: FC Reported: FC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FACS: Fluorescence-activated cell sorting; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

# **Properties**

Formulation: Phosphate-buffered saline, pH 7.2, containing 0.09% sodium azide and 0.1% gelatin

Purification: The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The

solution is free of unconjugated FITC.

Stability and Storage: Product stable at 2 - 8°C when stored undiluted. Do not freeze. Protect product from prolonged exposure to

light. For product expiry date, please contact techsupport@stemcell.com.

Directions for Use: For flow cytometry the suggested use of this antibody is 5 µL per 1 x 10^6 cells in 100 µL. It is recommended


that the antibody be titrated for optimal performance for each application.

#### Anti-Human CD45 Antibody, Clone 2D1, FITC

## **Antibodies**



### Data



Flow cytometry analysis of human peripheral blood mononuclear cells (PBMCs) labeled with Anti-Human CD45 Antibody, Clone 2D1, FITC (filled histogram) or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21, FITC (Catalog #60070FI) (solid line histogram).

### **Related Products**

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

### References

- 1. Barclay AN et al. (Eds.). (1997) The Leucocyte Antigen FactsBook, Second Edition. (pp. 244-7), NY: Academic Press.
- 2. Schwinzer R. (1989) In: Knapp W et al. (Eds.), Leucocyte Typing IV: White Cell Differentiation Antigens (pp. 628–34). New York, NY: Oxford University Press.
- 3. Shah VO et al. (1988) Flow cytometric analysis of human bone marrow. IV. Differential quantitative expression of T-200 common leukocyte antigen during normal hemopoiesis. J Immunol 140(6): 1861–7.
- 4. Dalchau R et al. (1980) Monoclonal antibody to a human leukocyte-specific membrane glycoprotein probably homologous to the leukocyte-common (L-C) antigen of the rat. Eur J Immunol 10(10): 737–44.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.