

**Goat Anti-Rabbit IgG (H+L)
Antibody, Polyclonal, AMCA**



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Goat polyclonal antibody against
rabbit IgG (H+L), AMCA-conjugated

Catalog #60129AV

1.5 mg

Product Description

AMCA-conjugated goat anti-rabbit IgG (H+L) antibody reacts with the heavy chains on rabbit IgG and the light chains common in most rabbit immunoglobulins. This antibody has minimal cross-reactivity to human, mouse, and rat serum proteins but may react to immunoglobulins from other species.

Target Antigen Name:	IgG (H+L), Rabbit
Alternative Names:	Not applicable
Gene ID:	Not applicable
Species Reactivity:	Rabbit IgG (H+L). Minimal cross reactivity to human, mouse, and rat serum proteins
Host Species:	Goat
Clonality:	Polyclonal
Clone:	Not applicable
Isotype:	Not applicable
Immunogen:	Not applicable
Conjugate:	AMCA

Applications

Verified:	ICC
Reported:	FC, ICC, IF, IHC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Freeze-dried from a solution containing sodium phosphate, sodium chloride, bovine serum albumin, and sodium azide
Purification:	The antibody was purified by antigen affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Protect product from prolonged exposure to light. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	Centrifuge vial before opening. Resuspend the product in 1.1 mL deionized water, this is the stock dilution. Centrifuge if solution is not clear. Prepare working dilution fresh each day. NOTE: Once resuspended, store stock dilution at 2 - 8°C and use within 6 weeks or aliquot and store at -80°C. Alternatively, add glycerol at 1:1 after resuspension and store as a liquid at -20°C. Avoid repeated freeze-thaw cycles. For immunocytochemistry the suggested concentration of this antibody is 1:100 dilution. It is recommended that the antibody be titrated for optimal performance for each application.

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Ferreira JS et al. (2015) GluN2B-containing NMDA receptors regulate AMPA receptor traffic through anchoring of the synaptic proteasome. *J Neurosci* 35(22): 8462–79. (ICC, IF)
2. Zimberlin CD et al. (2015) HDAC1 and HDAC2 collectively regulate intestinal stem cell homeostasis. *FASEB J* 29(5): 2070–80. (IF, IHC)
3. Fukumoto R et al. (2014) Ciprofloxacin enhances stress erythropoiesis in spleen and increases survival after whole-body irradiation combined with skin-wound trauma B. Ryffel, ed. *PLoS One* 9(2): e90448. (IF, IHC)
4. Posod A et al. (2014) The common antitussive agent dextromethorphan protects against hyperoxia-induced cell death in established in vivo and in vitro models of neonatal brain injury. *Neuroscience* 274: 260–72. (IHC, WB)
5. Schenke-Layland K et al. (2007) Collagen IV induces trophoectoderm differentiation of mouse embryonic stem cells. *Stem Cells* 25(6): 1529–38. (ICC, IF)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.