

**Anti-Human CD326 (EpCAM)
Antibody, Clone 5E11.3.1**



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Antibodies

Mouse monoclonal IgG1 antibody
against human CD326 (EpCAM),
unconjugated

Catalog #60147

100 µg 1 mg/mL

Product Description

The 5E11.3.1 antibody reacts with the human epithelial cell adhesion molecule (EpCAM or CD326), an ~40 kDa type I transmembrane glycoprotein, which functions as a homophilic and Ca⁺⁺-independent adhesion molecule. This antibody shows positive staining on luminal cells of normal human mammary epithelium. No staining to weak staining is observed within the basal cell compartment of the mammary epithelium. The 5E11.3.1 antibody gives variable staining on malignant breast epithelium. Positive staining is observed on the following cell lines: MCF7, CAMA, T47D, BT10, BT20, COLO 205, and WIDR. Staining is negative on human bone marrow cells, blood cells, lymphoid and mesenchymal tissues, and mammary and marrow fibroblast cultures. EpCAM has roles in several cellular processes, including signaling, migration, proliferation, and differentiation. It is expressed early during erythropoiesis and has been identified as a stem cell marker. The EpCAM protein is found on the basolateral membrane of most normal epithelial cells and is highly expressed in bone marrow, colon, and lung. EpCAM is not expressed on mesothelial cells or mesotheliomas, and so is widely employed as a diagnostic marker to distinguish mesothelioma and carcinoma cells, as well as serving as a target for immunotherapeutic treatment of carcinomas.

Target Antigen Name:	CD326 (EpCAM)
Alternative Names:	EGP2, EpCAM, Epithelial cell adhesion molecule, ESA, TACSTD1, TROP-1
Gene ID:	4072
Species Reactivity:	Human
Host Species:	Mouse (BALB/c)
Clonality:	Monoclonal
Clone:	5E11.3.1
Isotype:	IgG1
Immunogen:	Human mammary carcinoma cell line T47D
Conjugate:	Unconjugated

Applications


Verified:	FC
Reported:	FC, IF, IHC

Abbreviations: CellSep: Cell separation; ChIP: Chromatin immunoprecipitation; FA: Functional assay; FC: Flow cytometry; ICC: Immunocytochemistry; IF: Immunofluorescence microscopy; IHC: Immunohistochemistry; IP: Immunoprecipitation; RIA: Radioimmunoassay; WB: Western blotting

Properties

Formulation:	Phosphate-buffered saline
Purification:	The antibody was purified by affinity chromatography.
Stability and Storage:	Product stable at 2 - 8°C when stored undiluted. Do not freeze. For product expiry date, please contact techsupport@stemcell.com .
Directions for Use:	The suggested use of this antibody is: FC, 0.1 - 0.3 µg per 1 x 10 ⁶ cells in 100 µL volume; IHC, 1 - 3 µg/mL (a heat-induced antigen retrieval step should be used when working with formalin-fixed paraffin-embedded sections). It is recommended that the antibody be titrated for optimal performance for each application.

Data



Flow cytometry analysis of human MCF7 cells labeled with Anti-Human CD326 (EpCAM) Antibody, Clone 5E11.3.1, followed by Goat Anti-Mouse IgG Antibody, Polyclonal, FITC (Catalog #60138FI) (filled histogram), or Mouse IgG1, kappa Isotype Control Antibody, Clone MOPC-21 (Catalog #60070), followed by Goat Anti-Mouse IgG Antibody, Polyclonal, FITC (solid line histogram).

Related Products

For a complete list of antibodies, including other conjugates, sizes and clones, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/antibodies or contact us at techsupport@stemcell.com.

References

1. Aupet S et al. (2013) Isolation of viable human hepatic progenitors from adult livers is possible even after 48 hours of cold ischemia. *Tissue Eng Part C Methods* 19(7): 497–506. (IHC)
2. Kryczek I et al. (2009) Phenotype, distribution, generation, and functional and clinical relevance of Th17 cells in the human tumor environments. *Blood* 114(6): 1141–9. (FC, IHC)
3. Stingl J et al. (1998) Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast. *Differentiation* 63(4): 201–13. (FC, IF, IHC)
4. Bardy P et al. (1997) Isolation and analysis of different subpopulations of normal human breast epithelial cells after their infection with a retroviral vector encoding a cell surface marker. *Breast Cancer Res Treat* 44(2): 153–65. (FC)

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.