

Cytokines

Human Recombinant FGF-7 (KGF)

Fibroblast growth factor 7



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	78046	10 µg
	78046.1	50 µg
	78046.2	1000 µg

Product Description

Fibroblast growth factor 7 (FGF-7) is a member of the FGF family, and acts exclusively through a subset of FGF receptor isoforms expressed predominantly by epithelial cells (Finch & Rubin). FGF-7 seems to act specifically on epithelial cells and stimulates proliferation, migration, and differentiation of these cells, and also participates in epithelial protection and repair both in vitro and in vivo (Finch & Rubin; Werner). In contrast, FGF-7 is produced solely by cells of mesenchymal origin, and function as a paracrine mediator of mesenchymal-epithelial communication (Rubin et al.). FGF-7 has also been shown to supplement several wound-healing properties of bioengineered skin (Erdag et al.) and induce autophagy in human keratinocytes (Belleudi et al.). Additionally, FGF-7 has a role in pluripotent stem cell differentiation to endodermal pancreatic-like insulin producing cells and thymic epithelial cells (Inami et al.; Niu et al.).

Product Information

Alternative Names:	Fibroblast growth factor-7, HBGF-7, Heparin-binding growth factor 7, Keratinocyte growth factor
Accession Number:	P21781
Amino Acid Sequence:	MCNDMTPEQM ATNVNCSSPE RHTRSVDYME GGDIRVRLRF CRTQWYLRIK KRGKVKGTQE MKNNNYIMEI RTVAVGIVAI KGVSEFYLA MNKEGKLYAK KECNEDCNFK ELILENHYNT YASAKWTHNG GEMFVALNQK GIPVRGKGTK KEQKTAHFLP MAIT
Predicted Molecular Mass:	19 kDa
Species:	Human
Cross Reactivity:	Mouse, Rat
Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 5 \times 10^5$ units/mg ($EC_{50} \leq 2$ ng/mL) as determined by a cell proliferation assay of 4MBr-5 cells.
Purity:	≥ 95 %
Endotoxin Level:	Measured by kinetic limulus amoebocyte lysate (LAL) analysis and is ≤ 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Resuspend the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 2 weeks or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant FGF-7 (KGF) was tested by its ability to promote the proliferation of 4MBr-5 cells. Cell proliferation was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC₅₀ in the example above is less than 2 ng/mL.

(B) 2 µg of Human Recombinant FGF-7 (KGF) was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant FGF-7 (KGF) has a predicted molecular mass of 19 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Belleudi F et al. (2014) FGF7/KGF regulates autophagy in keratinocytes: A novel dual role in the induction of both assembly and turnover of autophagosomes. *Autophagy* 10(5): 803–21.
- Erdag G et al. (2004) FGF-7 expression enhances the performance of bioengineered skin. *Mol Ther* 10(1): 76–85.
- Finch PW & Rubin JS. (2004) Keratinocyte growth factor/fibroblast growth factor 7, a homeostatic factor with therapeutic potential for epithelial protection and repair. *Adv Cancer Res* 91: 69–136.
- Inami Y et al. (2011) Differentiation of induced pluripotent stem cells to thymic epithelial cells by phenotype. *Immunol Cell Biol* 89(2): 314–21.
- Niu J et al. (2007) Keratinocyte growth factor/fibroblast growth factor-7-regulated cell migration and invasion through activation of NF-kappaB transcription factors. *J Biol Chem* 282(9): 6001–11.
- Rubin JS et al. (1995) Keratinocyte growth factor. *Cell Biol Int* 19(5): 399–411.
- Werner S. (1998) Keratinocyte growth factor: a unique player in epithelial repair processes. *Cytokine Growth Factor Rev* 9(2): 153–65.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.