

Cytokines

Mouse Recombinant IGF-I



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Insulin-like growth factor 1

Catalog #	78078	10 µg
	78078.1	50 µg

Product Description

Insulin-like growth factor 1 (IGF-I) is a polypeptide which belongs to the family of insulin-like growth factors that are similar in molecular structure to proinsulin. IGF-I binds to the IGF-I receptor and is a potent activator of the PI3K/AKT pathway, it also activates ERK1/2 signalling. IGF-I is required for embryonic development, and it is produced mainly in liver in response to a hepatocyte growth hormone. In the absence of insulin, IGF-I is necessary for the maintenance of human pluripotent stem cells (Wang et al.). Together with interleukin 3 (IL-3), IGF-I stimulates differentiation and proliferation of myeloid cells, and has been shown to regulate lymphopoiesis by stimulating proliferation and differentiation of T and B cells in lymphoid organs (Heemskerk et al.).

Product Information

Alternative Names:	IBP1, IGF-IA, IGF-IB, IGF1A, Mechano growth factor, MGF, Somatomedin C
Accession Number:	P05017
Amino Acid Sequence:	MGPETLCGAE LVDALQFVCG PRGFYFNKPT GYGSSIRRAP QTGIVDECCF RSCDLRRLEM YCAPLKPTKA A
Predicted Molecular Mass:	7.8 kDa
Species:	Mouse
Cross Reactivity:	Human, Rat
Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 1 \times 10^5$ units/mg ($EC_{50} \leq 10$ ng/mL) as determined by a cell proliferation assay of FDC-P1 cells.
Purity:	$\geq 95\%$
Endotoxin Level:	Measured by kinetic limulus amebocyte lysate (LAL) analysis and is ≤ 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Resuspend the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 2 weeks or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Mouse Recombinant IGF-I was tested by its ability to promote the proliferation of FDC-P1 cells. Cell proliferation was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC₅₀ in the example above is less than 10 ng/mL.

(B) 2 µg of Mouse Recombinant IGF-I was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant IGF-I has a predicted molecular mass of 7.8 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Heemskerk VH et al. (1999) Insulin-like growth factor-1 (IGF-1) and growth hormone (GH) in immunity and inflammation. *Cytokine Growth Factor Rev* 10(1): 5–14.
- Wang L et al. (2007) Self-renewal of human embryonic stem cells requires insulin-like growth factor-1 receptor and ERBB2 receptor signaling. *Blood* 110(12): 4111–9.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.