Cytokines

Human Recombinant IL-6R

alpha

Interleukin 6 receptor alpha

Catalog # 78083

10 µg 50 µg

78083.1

Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Interleukin 6 receptor (IL-6R) alpha is a type I transmembrane glycoprotein that forms a complex with type I transmembrane signal transducer protein gp130 (CD130) and mediates the biological activities of IL-6. IL-6 binds to the membrane-bound non-signaling IL-6R alpha (mIL-6R), and the complex binds to two molecules of gp130 and leads to 'classical' IL-6-signal transduction, which includes activation of JAK/STAT, ERK, and PI3K signal transduction pathways (Scheller et al.). In contrast, a soluble form of IL-6R alpha (sIL-6R), which comprises the extracellular portion of the receptor, binds to the secreted IL-6 to form a complex that promotes bioavailability of IL-6. The complex of IL-6 and sIL-6R can bind to gp130 on cells that do not express the IL-6R and are unresponsive to IL-6. This process is known as trans-signaling (Hunter & Jones; Rose-John). sIL-6R regulates both local and systemic IL-6-mediated events. Elevated levels of sIL-6R have been documented in several disease conditions such as rheumatoid arthritis, myeloma, and Crohn's disease (Jones et al.; Mihara et al.).

Product Information

Alternative Names: B cell stimulatory factor-2, CD126, IL-6R, IL-6R1, IL-6RA, Interleukin-6 receptor, Interleukin-6 receptor alpha

Accession Number: P08887

Amino Acid Sequence: LAPRRCPAQE VARGVLTSLP GDSVTLTCPG VEPEDNATVH WVLRKPAAGS HPSRWAGMGR RLLLRSVQLH

> DSGNYSCYRA GRPAGTVHLL VDVPPEEPQL SCFRKSPLSN VVCEWGPRST PSLTTKAVLL VRKFQNSPAE DFQEPCQYSQ ESQKFSCQLA VPEGDSSFYI VSMCVASSVG SKFSKTQTFQ GCGILQPDPP ANITVTAVAR NPRWLSVTWQ DPHSWNSSFY RLRFELRYRA ERSKTFTTWM VKDLQHHCVI HDAWSGLRHV VQLRAQEEFG

QGEWSEWSPE AMGTPWTESR SPPAENEVST PMQALTTNKD DDNILFRDSA NATSLPVQDS SSVPLP

Predicted Molecular Mass: 38.6 kDa Species: Human Cross Reactivity: Mouse

Formulation: Lyophilized after dialysis against phosphate-buffered saline.

Source: CHO

Specifications

Activity: The specific activity is $\geq 5 \times 10^3$ units/mg (EC50 ≤ 200 ng/mL) as determined by the ability to inhibit

growth of M1 cells in the presence of 10 ng/mL human IL-6.

Purity:

Endotoxin Level: Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 0.2 EU/µg protein.

Preparation and Storage

Storage: Store at -80°C.

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Reconstitute the product in sterile water or phosphate-buffered saline to at


> least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. As a general guide, do not store at 2 - 8°C for more than 1 week or at -20°C for more than 2 months. Avoid repeated freeze-thaw cycles.

Human Recombinant IL-6R alpha

Cytokines



Data



(A) The biological activity of Human Recombinant IL-6R alpha was tested by its ability to inhibit the proliferation of M1 cells. Inhibition of cell proliferation was measured using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation inhibition is at 50% of maximum. The EC50 in the above example is less than 200 ng/mL.

(B) 5 μg of Human Recombinant IL-6R alpha was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant IL-6R alpha has a predicted molecular mass of 38.6 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

Hunter CA & Jones SA. (2015) IL-6 as a keystone cytokine in health and disease. Nat Immunol 16(5): 448–57. Jones SA et al. (2001) The soluble interleukin 6 receptor: mechanisms of production and implications in disease. FASEB J 15(1): 43–58. Mihara M et al. (2012) IL-6/IL-6 receptor system and its role in physiological and pathological conditions. Clin Sci (Lond) 122(4): 143–59. Rose-John S. (2012) IL-6 trans-signaling via the soluble IL-6 receptor: importance for the pro-inflammatory activities of IL-6. Int J Biol Sci 8(9): 1237–47.

Scheller J et al. (2011) The pro- and anti-inflammatory properties of the cytokine interleukin-6. Biochim Biophys Acta 1813(5): 878–88.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.