

Cytokines

Human Recombinant MCP-1 (CCL2)

Monocyte chemotactic protein 1



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 78087
78087.1

5 µg
20 µg

Product Description

Monocyte chemotactic protein-1 (MCP-1) also known as CCL2 is a member of CC family of chemokines. The protein is primarily induced by platelet-derived growth factor (PDGF) gene (Cochran et al.). The biological effects of MCP-1 are mediated via specific G-protein-coupled receptor CCR2, which in turn activates signal transduction pathways leading to monocyte transmigration (Sozzani et al.). Migration of monocytes from the bloodstream across the vascular endothelium is required for routine immunological surveillance of tissues, as well as other immunomodulatory effects. MCP-1 is produced by a variety of cell types, including endothelial, fibroblasts, epithelial, smooth muscle, mesangial, astrocytic, monocytic, and microglial cells, which are important for antiviral responses in the peripheral circulations and in tissues (Cushing et al.; Deshmane et al.). MCP-1 play a role in physiological processes such as neurogenesis, neuroprotection, and neurotransmission and has important implications in neurological disorders like multiple sclerosis and alzheimer's disease, where it is produced during neuroinflammation at the sites of lesions (Conductier et al.).

Product Information

Alternative Names: DIA, Differentiation-inducing factor, Leukemia inhibitory factor, HILDA, Human interleukin in DA cells, MCAF, Melanoma-derived LPL inhibitor, MLPLI, Monocyte chemoattractant protein 1

Accession Number: P13500

Amino Acid Sequence: QPDAINAPVT CCYNFTNRKI SVQRLASYRR ITSSKCPKEA VIFKTIVAKE ICADPKQKWV QDSMDHLDKQ TQTPKT

Predicted Molecular Mass: 8.7 kDa

Species: Human

Cross Reactivity: Mouse

Formulation: Lyophilized from a sterile filtered aqueous solution containing 0.1% trifluoroacetic acid.

Source: E. coli

Specifications

Activity: Biological activity was detectable at ≤ 100 ng/mL as determined by a cell migration assay of THP-1 cells.

Purity: ≥ 95 %

Endotoxin Level: Measured by kinetic limulus ameocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.


Preparation and Storage

Storage: Store at -20°C to -80°C.

Stability: Stable as supplied for 12 months from date of receipt.

Preparation: Centrifuge vial before opening. Resuspend the product in sterile water containing 0.1% bovine serum albumin (BSA) to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 month or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles. NOTE: If reconstituted product will be used immediately BSA is not required.

Data



(A) The biological activity of Human Recombinant MCP-1 was tested by its ability to induce chemotaxis of THP-1 cells. Cell migration was measured after 1 hour using a fluorometric assay method. Increase in migration over basal level was seen starting at 1 ng/mL.

(B) 1 µg of Human Recombinant MCP-1 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant MCP-1 has a predicted molecular mass of 8.7 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Cochran BH et al. (1983) Molecular cloning of gene sequences regulated by platelet-derived growth factor. *Cell* 33(3): 939–47.
- Conductier G et al. (2010) The role of monocyte chemoattractant protein MCP1/CCL2 in neuroinflammatory diseases. *J Neuroimmunol* 224(1-2): 93–100.
- Cushing SD et al. (1990) Minimally modified low density lipoprotein induces monocyte chemotactic protein 1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci U S A* 87(13): 5134–8.
- Deshmane SL et al. (2009) Monocyte chemoattractant protein-1 (MCP-1): an overview. *J Interferon Cytokine Res* 29(6): 313–26.
- Sozzani S et al. (1994) Receptors and transduction pathways for monocyte chemotactic protein-2 and monocyte chemotactic protein-3. Similarities and differences with MCP-1. *J Immunol* 152(7): 3615–22.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.