

Cytokines

Human Recombinant Oncostatin M

Oncostatin M



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog #	78094	10 µg
	78094.1	50 µg
	78094.2	1000 µg

Product Description

Oncostatin M (OSM) is a member of interleukin 6 (IL-6) family of cytokines and bears close resemblance to leukemia-inhibitory factor (LIF) and granulocyte colony-stimulating factor (G-CSF) in amino acid sequence and its modulation of differentiation in a variety of cell types (Rose & Bruce). OSM signals through type I receptor (consisting of gp130 and LIF receptor (LIFR)) and type II receptor (consisting of gp130 and OSM receptor (OSMR)), which eventually activate the JAK/STAT pathway (Auguste et al.; Gómez-Lechón). OSM is primarily produced by activated T cells and monocytes, and also by activated macrophages, neutrophils, mast cells, and dendritic cells. OSM is also produced within the bone microenvironment by cells of both hematopoietic and mesenchymal origin including osteocytes and osteoblasts. OSM is involved in differentiation, cell proliferation, hematopoiesis, and inflammation, and also has been shown to have implications in liver development, bone formation and resorption (Sims & Quinn; Tanaka & Miyajima).

Product Information

Alternative Names:	OSM
Accession Number:	P13725
Amino Acid Sequence:	MAAIGSCSKE YRVLLGQLQK QTDLMQDTSR LLDPIYRIQG LDVPKLRHC RERPGAFPSE ETLRGLGRRG FLQTLNATLG CVLHRLADLE QRLPKAQDLE RSGLNIEDLE KLQMARPNIIL GLRNNIYCMA QLLDNSDTAE PTKAGRGASQ PPTPTPASDA FQRKLEGCRF LHGYHRFMHS VGRVFSKWGE SPNRSRRHSP HQALRKGVRP TRPSRKGKRL MTRGQLPR
Predicted Molecular Mass:	25.9 kDa
Species:	Human
Cross Reactivity:	Mouse
Formulation:	Lyophilized after dialysis against phosphate-buffered saline.
Source:	E. coli


Specifications

Activity:	The specific activity is $\geq 1 \times 10^5$ units/mg ($EC_{50} \leq 10$ ng/mL) as determined by a cell proliferation assay using TF-1 cells.
Purity:	≥ 95 %
Endotoxin Level:	Measured by kinetic Limulus amebocyte lysate (LAL) analysis and is ≤ 0.2 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Resuspend the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 2 weeks or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant Oncostatin M was tested by its ability to promote the proliferation of TF-1 cells. Cell proliferation was measured using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC50 in the above example is less than 10 ng/mL.

(B) 2 µg of Human Recombinant Oncostatin M was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant Oncostatin M has a predicted molecular mass of 25.9 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Auguste P et al. (1997) Signaling of type II oncostatin M receptor. *J Biol Chem* 272(25): 15760–4.
- Gómez-Lechón MJ. (1999) Oncostatin M: signal transduction and biological activity. *Life Sci* 65(20): 2019–30.
- Rose TM & Bruce AG. (1991) Oncostatin M is a member of a cytokine family that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. *Proc Natl Acad Sci U S A* 88(19): 8641–5.
- Sims NA & Quinn JMW. (2014) Osteoimmunology: oncostatin M as a pleiotropic regulator of bone formation and resorption in health and disease. *Bonekey Rep* 3: 527.
- Tanaka M & Miyajima A. (2003) Oncostatin M, a multifunctional cytokine. *Rev Physiol Biochem Pharmacol* 149: 39–52.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.