

Cytokines

Mouse Recombinant VEGF-164



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Vascular endothelial growth factor

Catalog #	78102	20 µg
	78102.1	100 µg
	78102.2	1000 µg

Product Description

Vascular endothelial growth factor (VEGF) is a heparin-binding homodimeric glycoprotein involved in vasculogenesis and angiogenesis. VEGF binds to FLT1 (VEGFR-1) and KDR (VEGFR-2), and activates Raf/MEK/ERK and PI3K/AKT pathways (Ferrara et al.). VEGF exists in multiple isoforms that result from alternative splicing of VEGF mRNA in the terminal exon. Proximal splice-site selection in exon 8 results in pro-angiogenic VEGF_{xxx} isoforms (xxx is the number of amino acids), whereas distal splice-site selection results in anti-angiogenic VEGF_{xxx}b isoforms (Nowak et al.). VEGF plays an important role in neurogenesis both in vitro and in vivo (Storkebaum et al.). It has neurotrophic effects on neurons of the central nervous system, and it promotes growth and survival of dopaminergic neurons and astrocytes. VEGF also promotes growth and survival of vascular endothelial cells, monocyte chemotaxis, and colony formation by granulocyte-macrophage progenitor cells (Ferrara et al.). Various splice variants of VEGF exist, with different functions. For example, it has been shown that VEGF isoform VEGF-164(165) and not VEGF-120(121) induces inflammation, stimulates intracellular adhesion molecule (ICAM)-1 expression on endothelial cells, and induces chemotaxis of monocytes (Usui et al.).

Product Information

Alternative Names:	MGC70609, MVCD-1, Vascular endothelial growth factor 2, Vascular endothelial growth factor A, Vascular permeability factor, Vasculotropin, VEGFA, VPF
Accession Number:	Q00731-1
Amino Acid Sequence:	MAPTTEGEQK SHEVIKFMDV YQRSYCRPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCAGC CNDEALECVP TSESNITMQI MRIKPHQSQH IGEMSFLQHS RCECRPKKDR TKPEKHCEPC SERRKHLFVQ DPQCKCCK NTDSRCKARQ LELNERTCRC DKPRR
Predicted Molecular Mass:	39 kDa
Species:	Mouse
Cross Reactivity:	Human, Rat
Formulation:	Lyophilized after dialysis against HEPES and sodium chloride, pH 7.0.
Source:	P. pastoris

Specifications


Activity:	The specific activity is $\geq 1.7 \times 10^5$ units/mg ($EC_{50} \leq 6$ ng/mL) as determined by a cell proliferation assay using HUVECs.
Purity:	$\geq 97\%$
Endotoxin Level:	Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage:	Store at -80°C.
Stability:	Stable as supplied for 12 months from date of receipt.
Preparation:	Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -20°C for more than 6 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Mouse Recombinant VEGF-164 was tested by its ability to promote the proliferation of HUVECs. Cell proliferation was measured using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell proliferation is at 50% of maximum. The EC₅₀ in the above example is less than 6 ng/mL.

(B) 2 µg of Mouse Recombinant VEGF-164 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Mouse Recombinant VEGF-164 has a predicted molecular mass of 39 kDa.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Ferrara N et al. (2003) The biology of VEGF and its receptors. *Nat Med* 9(6): 669–76.
- Nowak DG et al. (2008) Expression of pro- and anti-angiogenic isoforms of VEGF is differentially regulated by splicing and growth factors. *J Cell Sci* 121(Pt 20): 3487–95.
- Storkebaum E et al. (2004) VEGF: once regarded as a specific angiogenic factor, now implicated in neuroprotection. *Bioessays* 26(9): 943–54.
- Usui T et al. (2004) VEGF164(165) as the pathological isoform: differential leukocyte and endothelial responses through VEGFR1 and VEGFR2. *Invest Ophthalmol Vis Sci* 45(2): 368–74.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.