

Dissociation Reagents

Collagenase Type V

For digestion of native collagen fibrils



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 07430
07431

100 mg
1 g

Product Description

Collagenase is a protease consisting of a single polypeptide chain approximately 1,000 amino acid residues in length. Collagenase is capable of digesting native collagen fibrils commonly found in connective tissues and therefore is frequently used for tissue dissociation. Collagenase preparations contain the activity of several proteases, including collagenase, caseinase, clostripain, and trypsin (Kessler & Yaron). Collagenase Type V contains high levels of collagenase and caseinase activities.

Product Information

Alternative Names:	Clostridium histolyticum collagenase; Collagenase 5; Collagenase Type 5; Collagenase V
Format:	Lyophilized powder
Storage:	Store at 2 - 8°C.
Stability:	Stable until expiry date (EXP) on label.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	68 - 130 kDa
CAS Number:	9001-12-1
Optimum pH:	6.3 - 8.5
Cleavage Site:	-Pro-X-†-Gly-Pro-Y- : X = neutral Y = nonspecific

Specifications

Source:	Clostridium histolyticum
Activity:	Collagenase: ≥ 450 CDU/mg dry weight (mgdw); Caseinase: ≥ 450 u/mgdw; Clostripain: ≤ 3.0 u/mgdw; Trypsin: ≤ 0.3 u/mgdw. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

Collagenase: 1 collagenase digestion unit (CDU) equals 1 μmol of L-leucine equivalents released from collagen in 5 hours at 37°C, pH 7.5.

Caseinase: 1 unit equals 1 μmol of L-leucine equivalents released from 25 mg vitamin-free casein in 5 hours at 37°C, pH 7.5. Measures non-specific proteolytic activity.

Clostripain: 1 unit hydrolyzes 1 μmol of N α -benzoyl-L-arginine ethyl ester (BAEE)/minute at 25°C at pH 7.6, after activation in 2.5 mM dithiothreitol (DTT).

Trypsin: 1 unit hydrolyzes 1 μmol of BAEE/minute at 25°C at pH 7.6.

References

Barrett AJ & Starkey PM. (1973) The Interaction of α 2-Macroglobulin with Proteinases. J Biochem 133: 709–24.

Kessler E & Yaron A. (1973) A novel aminopeptidase from clostridium histolyticum. Biochem Biophys Res Commun 50(2): 405–12.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2016 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design and Scientists Helping Scientists are trademarks of STEMCELL Technologies Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.