

Dissociation Reagents

Hyaluronidase

For hydrolysis of hyaluronic acid



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 07461
07462


50,000 Units
300,000 Units

Product Description

Hyaluronidase is a polysaccharidase consisting of a single polypeptide chain of 450 amino acid residues and contains four disulfide bonds. Hyaluronidase is glycosylated and contains 5% mannose and 2.17% glucosamine (Borders & Raftery). Hyaluronidase cleaves endo-N-acetylhexosaminic bonds in hyaluronic acid and chondroitin sulfate A and C to tetrasaccharide residues, as hyaluronic acid and chondroitin sulfate are often found in connective tissues. Hyaluronidase is frequently used with other proteases such as collagenase.

Product Information

Alternative Names:	Chondroitinase; Hyaluronate 4-glycanohydrolase; Hyaluronoglucosaminidase
Format:	Lyophilized powder
Storage:	Store at -20°C.
Stability:	Stable until expiry date (EXP) on label.
Reconstitution:	Dissociation reagents can be reconstituted in a balanced salt solution or buffer of choice.
Molecular Weight:	55 - 61 kDa
CAS Number:	37326-33-3
Optimum pH:	4.5 - 6.0
Cleavage Site:	Hyaluronidase randomly cleaves the 1,4-linkages between 2-acetamido-2-deoxy-b-D-glucose and D-glucose residues in hyaluronate.



Cleavage site of Hyaluronidase

Specifications

Source:	Bovine testes
Activity:	≥ 300 units/mg dry weight. See Notes for further information.

Related Products

For a complete list of dissociation reagents, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com or contact us at techsupport@stemcell.com.

Notes

ACTIVITY UNITS

1 unit is based on the change in absorbency (turbidity) at 540 nm of an internal standard assayed concurrently with each lot.

References

- Anway MD et al. (2002) Isolation of sertoli cells from adult rat testes: an approach to ex vivo studies of sertoli cell function. *Biol Reprod* 68(3): 996–1002.
- Babona-Pilipos R et al. (2012) A galvanotaxis assay for analysis of neural precursor cell migration kinetics in an externally applied direct current electric field. *J Vis Exp* (68): 1–8.
- Borders CL & Raftery MA. (1968) Purification and partial characterization of testicular hyaluronidase. *J Biol Chem* 243(13): 3756–62.
- Farias VA et al. (2011) Human umbilical cord stromal stem cell express CD10 and exert contractile properties. *Placenta* 32(1): 86–95.
- Folkman J et al. (1979) Long-term culture of capillary endothelial cells. *Proc Natl Acad Sci USA* 76(10): 5217–21.
- Gritti A et al. (1999) Epidermal and fibroblast growth factors behave as mitogenic regulators for a single multipotent stem cell-like population from the subventricular region of the adult mouse forebrain. *J Neurosci* 19(9): 3287–97.
- Ishimaru N et al. (1999) Estrogen deficiency accelerates autoimmune exocrinopathy in murine Sjögren's syndrome through fas-mediated apoptosis. *Am J Pathol* 155(1): 173–81.
- Labarge MA et al. (2013) Processing of human reduction mammoplasty and mastectomy tissues for cell culture. *J Vis Exp* (71): 1–7.
- Liagre B et al. (2004) Diosgenin, a plant steroid, induces apoptosis in human rheumatoid arthritis synoviocytes with cyclooxygenase-2 overexpression. *Arthritis Res Ther* 6(4): R373–83.
- Liu X et al. (2013) Nonlinear growth kinetics of breast cancer stem cells: implications for cancer stem cell targeted therapy. *Sci Rep* 3: 2473.
- Sauvageot CM-E et al. (2009) Efficacy of the HSP90 inhibitor 17-AAG in human glioma cell lines and tumorigenic glioma stem cells. *Neuro Oncol* 11(2): 109–21.
- Taylor RA et al. (2012) Human epithelial basal cells are cells of origin of prostate cancer, independent of CD133 status. *Stem Cells* 30(6): 1087–96.
- Tomooka Y et al. (1983) Growth effect of lithium on mouse mammary epithelial cells in serum-free collagen gel culture. *J Cell Physiol* 117(3): 290–6.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.