

Cytokines

Human Recombinant GDF-5



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 78111
78111.1

50 µg
100 µg

Product Description

Growth differentiation factor 5 (GDF-5) is a member of the bone morphogenic protein (BMP) subclass of the transforming growth factor beta (TGF-β) superfamily. It binds a receptor complex comprising BMPRI1B and BMPRI2, which then activates the SMAD signaling pathway (Carreira et al.; Nishitoh et al.; Osório et al.). GDF-5 enhances chondrogenic differentiation of mesenchymal stem cells, skeletogenesis, and dendrite growth during development (Coleman et al.; Francis-West et al.). Studies in rat demonstrated that GDF-5 regulates patterning, neurogenesis, astrogliogenesis, and neuronal specification (Gajavelli et al.; Kriegstein et al.; O'Keeffe et al.).

Product Information

Alternative Names: BMP-14, Bone morphogenic protein 14, CDMF-1, Growth differentiation factor 5
Accession Number: P43026
Amino Acid Sequence: MAPLATRQGK RPSKNLKARC SRKALHVNFK DMGWDDWIIA PLEYEAFHCE GLCEFPLRSH LEPTNHAIVQ TLMNSMDPES TPPTCCVPTR LSPISILFIDSA NNVVYKQY EDMVVESCGC R
Predicted Molecular Mass: 13.7 kDa monomer; 27.4 kDa dimer
Species: Human
Cross Reactivity: Mouse
Formulation: Lyophilized from a sterile-filtered aqueous solution containing 0.1% trifluoroacetic acid.
Source: E. coli

Specifications


Activity: The specific activity is $\geq 8.3 \times 10^2$ units/mg ($EC_{50} \leq 1.2 \mu\text{g/mL}$) as determined by alkaline phosphatase activity induced in ATDC-5 cells.
Purity: $\geq 95\%$
Endotoxin Level: Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1 EU/µg protein.

Preparation and Storage

Storage: Store at -20°C to -80°C.
Stability: Stable as supplied for 12 months from date of receipt.
Preparation: Centrifuge vial before opening. Reconstitute the product in sterile water to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Data



(A) The biological activity of Human Recombinant GDF-5 was tested by its ability to induce alkaline phosphatase production in ATDC-5 cells. Alkaline phosphatase production was measured after 63 hours using a fluorometric assay method. The EC50 is defined as the effective concentration of the growth factor at which alkaline phosphatase activity is at 50% of maximum. The EC50 in the above example is 0.017 μg/mL.

(B) 1 μg of Human Recombinant GDF-5 was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant GDF-5 has a predicted molecular mass of 27.4 kDa (13.7 kDa per monomer).

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Carreira AC et al. (2014) Bone morphogenetic proteins: facts, challenges, and future perspectives. *J Dent Res* 93(4): 335–45.
- Coleman CM et al. (2013) Growth differentiation factor-5 enhances in vitro mesenchymal stromal cell chondrogenesis and hypertrophy. *Stem Cells Dev* 22(13): 1968–76.
- Francis-West PH et al. (1999) Mechanisms of GDF-5 action during skeletal development. *Development* 126(6).
- Gajavelli S et al. (2004) BMP signaling initiates a neural crest differentiation program in embryonic rat CNS stem cells. *Exp Neurol* 188(2): 205–23.
- Kriegstein K et al. (1995) Trophic and protective effects of growth/differentiation factor 5, a member of the transforming growth factor-β superfamily, on midbrain dopaminergic neurons. *J Neurosci Res* 42(5): 724–32.
- Nishitoh H et al. (1996) Identification of type I and type II serine/threonine kinase receptors for growth/differentiation factor-5. *J Biol Chem* 271(35): 21345–52.
- O'Keeffe GW et al. (2004) Expression of growth differentiation factor-5 in the developing and adult rat brain. *Dev Brain Res* 151(1–2): 199–202.
- Osório C et al. (2013) Growth differentiation factor 5 is a key physiological regulator of dendrite growth during development. *Development* 140(23).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2018 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.