

Cytokines

Human Recombinant Myostatin

Myostatin



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 78119
78119.1

10 µg
100 µg

Product Description

Myostatin is a member of transforming growth factor beta (TGF-β) superfamily that signals by binding to activin type II receptor, resulting in the recruitment of either ALK3 or ALK4 coreceptor, and activation of SMAD2/3 (Kondás et al.). It is released by myocytes and inhibits muscle growth and ability to regenerate (Lee & Lee). Myostatin appears to affect the lipid catabolic metabolism of adipocytes and inhibits preadipocyte differentiation, as demonstrated in the 3T3-L1 cell line (Li et al.).

Product Information

Alternative Names: GDF-8, Growth differentiation factor 8, MSTN78
Accession Number: O14793
Amino Acid Sequence: DFGLDCDEHS TESRCCRYPL TVDFEAFGWD WIIAPKRYKA NYCSGECEFV FLQKYPHTHL VHQANPRGSA GPCCTPTKMS PINMLYFNGK EQIIYGKIPA MVVDRCGCS
Predicted Molecular Mass: 12.4 kDa monomer, 24.8 kDa dimer
Species: Human
Cross Reactivity: Mouse, Rat
Formulation: Lyophilized from a sterile filtered aqueous solution containing 0.1% trifluoroacetic acid.
Source: E. coli


Specifications

Activity: The specific activity is $\geq 2.0 \times 10^4$ units/mg ($EC_{50} \leq 50$ ng/mL) as determined by cytotoxicity assay of MPC-11 cells.
Purity: $\geq 95\%$
Endotoxin Level: Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is ≤ 1.0 EU/µg protein.

Preparation and Storage

Storage: Store at -20°C to -80°C.
Stability: Stable as supplied for 12 months from date of receipt.
Preparation: Centrifuge vial before opening. Resuspend the product in 20 mM hydrochloric acid containing 0.1% bovine serum albumin (BSA) to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex. Store at 2 - 8°C for up to 1 month or at -20°C to -80°C for up to 3 months. Avoid repeated freeze-thaw cycles. NOTE: If reconstituted product will be used immediately BSA is not required.

Data



(A) The biological activity of Human Recombinant Myostatin was tested by its ability to inhibit proliferation of MPC-11 cells. Cell viability was measured after 66 hours using a fluorometric assay method. The EC₅₀ is defined as the effective concentration of the growth factor at which cell viability is at 50% of maximum. The EC₅₀ in the above example is 12 ng/mL.

(B) 1 µg of Human Recombinant Myostatin was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining.

Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, please visit our website at www.stemcell.com/cytokines or contact us at techsupport@stemcell.com.

References

- Kondás K et al. (2008) Both WFIKKN1 and WFIKKN2 have high affinity for growth and differentiation factors 8 and 11. *J Biol Chem* 283(35): 23677–84.
- Lee Y-S & Lee S-J. (2013) Regulation of GDF-11 and myostatin activity by GASP-1 and GASP-2. *Proc Natl Acad Sci USA* 110(39): E3713–22.
- Li F et al. (2011) Myostatin regulates preadipocyte differentiation and lipid metabolism of adipocyte via ERK1/2. *Cell Biol Int* 35(11): 1141–6.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.