Dibutyryl-cAMP

Small Molecules

cAMP pathway activator; Activates cAMP-dependent protein kinases

Catalog # 73882 25 mg 73884 100 mg 73886 250 mg



Scientists Helping Scientists™ | www.stemcell.com

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713 INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

Dibutyryl-cAMP is a cell-permeable cyclic AMP (cAMP) analog that activates cAMP-dependent protein kinases (Schwede et al.). This product is supplied as the sodium salt of the molecule.

Molecular Name: Dibutyryl-cAMP (Sodium Salt)

Alternative Names: Bucladesine; DC 2797

Purity: $\geq 95\%$

Chemical Name: N-(1-oxobutyl)-cyclic 3',5'-(hydrogen phosphate) 2'-butanoate-adenosine, monosodium salt

Structure:

Properties

Physical Appearance: A crystalline solid

Storage: Product stable at -20°C as supplied. Protect product from prolonged exposure to light. For long-term storage

store with a desiccant.

Stable as supplied for 12 months from date of receipt.

Solubility: $\cdot PBS \le 6.5 \text{ mM}$

 \cdot DMSO ≤ 6.5 mM

· Absolute ethanol ≤ 2 mM

For example, to prepare a 5 mM stock solution in DMSO, resuspend 10 mg in 4.07 mL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Small Molecules Dibutyryl-cAMP



Published Applications

DIFFERENTIATION

- · Suppresses experimental autoimmune encephalomyelitis development by reducing demyelination and mobilizing neural stem cells in the subventricular zone toward the demyelinated plaques (Khezri et al.).
- · Induces intrinsic axon growth in peripheral and central nervous systems, and morphological differentiation of astrocytes (Imamura & Ozawa; Knott et al.).
- · Stimulates neurite outgrowth in PC12 cells (Maruoka et al.).

References

Imamura M & Ozawa E. (1998) Differential expression of dystrophin isoforms and utrophin during dibutyryl-cAMP-induced morphological differentiation of rat brain astrocytes. Proc Natl Acad Sci USA 95(11): 6139–44.

Khezri S et al. (2013) Dibutyryl cyclic AMP inhibits the progression of experimental autoimmune encephalomyelitis and potentiates recruitment of endogenous neural stem cells. J Mol Neurosci 51(2): 298–306.

Knott EP et al. (2014) Cyclic AMP signaling: a molecular determinant of peripheral nerve regeneration. Biomed Res Int 2014: 651625. Maruoka H et al. (2010) Dibutyryl-cAMP up-regulates nur77 expression via histone modification during neurite outgrowth in PC12 cells. J Biochem 148(1): 93–101.

Schwede F et al. (2000) Cyclic nucleotide analogs as biochemical tools and prospective drugs. Pharmacol Ther 87(2-3): 199-226.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, visit www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.