

## Small Molecules

### Troglitazone

NF- $\kappa$ B pathway inhibitor; Activates PPAR $\gamma$

Catalog # 73892  
73894

5 mg  
10 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM


FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

## Product Description

Troglitazone is a potent and selective activator of peroxisome proliferator-activated receptor- $\gamma$  (PPAR $\gamma$ ) with EC50 = 0.55 and 0.78  $\mu$ M for human and mouse, respectively. Troglitazone does not inhibit PPAR $\alpha$  or PPAR $\delta$  at up to 10  $\mu$ M (Willson et al.). Troglitazone causes cell cycle arrest at phase G1, inducing caspase-dependent apoptosis in both HeLa and hepatocellular carcinoma cell lines (Yoshizawa et al.; Chang et al.).

Molecular Name:	Troglitazone
Alternative Names:	Resulin; Rezulin
CAS Number:	97322-87-7
Chemical Formula:	C <sub>24</sub> H <sub>27</sub> NSO <sub>5</sub>
Molecular Weight:	441.5 g/mol
Purity:	≥ 98%
Chemical Name:	5-[[4-[(3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione

Structure:



## Properties

Physical Appearance:	A crystalline solid
Storage:	Product stable at -20°C as supplied. Protect product from prolonged exposure to light. For long-term storage store with a desiccant. For product expiry date, please contact techsupport@stemcell.com.
Solubility:	· DMSO ≤ 65 mM · Absolute ethanol ≤ 0.5 mM For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 227 $\mu$ L of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

Compound has low solubility in aqueous media. For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

## Published Applications

### DIFFERENTIATION

- Stimulates adipogenesis in 3T3-L1 cells (Jeong & Yoon).
- Inhibits macrophage differentiation (Chen Y et al.).

### CANCER RESEARCH

- Inhibits growth and induces apoptosis in non-small cell lung carcinoma, bladder cancer, cervical cancer, and prostate cancer cells (Chen H-M et al.; Santha et al.; Satoh et al.; Yan et al.).
- Promotes cytostatic effects in MDA-MB-231 and MCF-7 breast cancer cell lines (Berthe et al.).

## References

- Berthe A et al. (2016)  $\Delta 2$ -Troglitazone promotes cytostatic rather than pro-apoptotic effects in breast cancer cells cultured in high serum conditions. *Cell Cycle* 15(24): 3402–12.
- Chang HK et al. (2017) Inhibition of ERK activity enhances the cytotoxic effect of peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) agonists in HeLa cells. *Biochem Biophys Res Commun* 482(4): 843–8.
- Chen H-M et al. (2014) Ubiquitination of p53 is involved in troglitazone induced apoptosis in cervical cancer cells. *Asian Pac J Cancer Prev* 15(5): 2313–8.
- Chen Y et al. (2014) The inhibitory effect of troglitazone on macrophage differentiation mediated by repressing NF- $\kappa$ B activation independently of PPAR $\gamma$ . *Mol Cell Toxicol* 10(3): 261–8.
- Jeong S & Yoon M. (2011)  $17\beta$ -Estradiol inhibition of PPAR $\gamma$ -induced adipogenesis and adipocyte-specific gene expression. *Acta Pharmacol Sin* 32(2): 230–8.
- Lee MK & Olefsky JM. (1995) Acute effects of troglitazone on in vivo insulin action in normal rats. *Metabolism* 44(9): 1166–9.
- Santha S et al. (2015) Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-troglitazone-induced apoptosis in prostate cancer cells involve AMP-activated protein kinase. *J Biol Chem* 290(36): 21865–75.
- Satoh T et al. (2002) Activation of peroxisome proliferator-activated receptor- $\gamma$  stimulates the growth arrest and DNA-damage inducible 153 gene in non-small cell lung carcinoma cells. *Oncogene* 21(14): 2171–80.
- Willson TM et al. (2000) The PPARs: from orphan receptors to drug discovery. *J Med Chem* 43(4): 527–50.
- Yan S et al. (2014) The PPAR $\gamma$  agonist Troglitazone induces autophagy, apoptosis and necroptosis in bladder cancer cells. *Cancer Gene Ther* 21(5): 188–93.
- Yoshizawa K et al. (2002) Peroxisome proliferator-activated receptor gamma ligand troglitazone induces cell cycle arrest and apoptosis of hepatocellular carcinoma cell lines. *Cancer* 95(10): 2243–51.

## Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at [www.stemcell.com/smallmolecules](http://www.stemcell.com/smallmolecules) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.