

Small Molecules

BMS 493

Retinoid pathway activator; Activates pan-retinoic acid receptor

Catalog # 73972
73974

1 mg
10 mg



Scientists Helping Scientists™ | WWW.STEMCELL.COM

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713


INFO@STEMCELL.COM • TECHSUPPORT@STEMCELL.COM

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Product Description

BMS 493 is an inverse agonist of the pan-retinoic acid receptors (RARs). RARs are ligand-dependent transcription factors that regulate gene networks that control cell growth, differentiation, survival, and death (Germain et al.). BMS 493 has been shown to increase nuclear receptor co-repressor (NCoR) interaction with RARs (Germain et al.; Rinkevich et al.).

Molecular Name: BMS 493
Alternative Names: BMS 204493
CAS Number: 215030-90-3
Chemical Formula: C₂₉H₂₄O₂
Molecular Weight: 404.5 g/mol
Purity: ≥ 98%
Chemical Name: 4-[(E)-2-[5,5-dimethyl-8-(2-phenylethynyl)-6H-naphthalen-2-yl]ethenyl]benzoic acid
Structure:



Properties

Physical Appearance: A crystalline solid
Storage: Product stable at -20°C as supplied. Protect product from prolonged exposure to light. For long-term storage store with a desiccant. For product expiry date, please contact techsupport@stemcell.com.
Solubility: · DMSO ≤ 60 mM
For example, to prepare a 10 mM stock solution in DMSO, resuspend 1 mg in 247 µL of DMSO.

Prepare stock solution fresh before use. Information regarding stability of small molecules in solution has rarely been reported, however, as a general guide we recommend storage in DMSO at -20°C. Aliquot into working volumes to avoid repeated freeze-thaw cycles. The effect of storage of stock solution on compound performance should be tested for each application.

For use as a cell culture supplement, stock solution should be diluted into culture medium immediately before use. Avoid final DMSO concentration above 0.1% due to potential cell toxicity.

Published Applications

DIFFERENTIATION

- Affects development of bronchial tubule formation in mice (Chazaud et al.).
- Inhibits activation of immature dendritic cells (Geissmann et al.).

CANCER RESEARCH

- Inhibits differentiation of leukemic blasts (Kamashev et al.).

References

- Chazaud C et al. (2003) Retinoic acid signaling regulates murine bronchial tubule formation. *Mech Dev* 120(6): 691–700.
- Geissmann F et al. (2003) Retinoids regulate survival and antigen presentation by immature dendritic cells. *J Exp Med* 198(4): 623–34.
- Germain P et al. (2009) Differential action on coregulator interaction defines inverse retinoid agonists and neutral antagonists. *Chem Biol* 16(5): 479–89.
- Kamashev D et al. (2004) PML-RARA-RXR oligomers mediate retinoid and rexinoid/cAMP cross-talk in acute promyelocytic leukemia cell differentiation. *J Exp Med* 199(8): 1163–74.
- Rinkevich Y et al. (2007) Systemic bud induction and retinoic acid signaling underlie whole body regeneration in the urochordate *Botryllodes leachi*. *PLoS Biol* 5(4): e71.

Related Small Molecules

For a complete list of small molecules available from STEMCELL Technologies, please visit our website at www.stemcell.com/smallmolecules or contact us at techsupport@stemcell.com.

This product is hazardous. Please refer to the Safety Data Sheet (SDS).

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.