

## Cytokines

### Human Recombinant GDF-11, ACF

Growth differentiation factor 11,  
animal component-free



Scientists Helping Scientists™ | [WWW.STEMCELL.COM](http://WWW.STEMCELL.COM)

TOLL FREE PHONE 1 800 667 0322 • PHONE +1 604 877 0713

[INFO@STEMCELL.COM](mailto:INFO@STEMCELL.COM) • [TECHSUPPORT@STEMCELL.COM](mailto:TECHSUPPORT@STEMCELL.COM)

FOR GLOBAL CONTACT DETAILS VISIT OUR WEBSITE

Catalog # 78189  
78189.1  
78189.2

20 µg  
100 µg  
1000 µg

## Product Description

Growth differentiation factor 11 (GDF-11) is a member of transforming growth factor beta (TGF- $\beta$ ) family. It binds to the TGF- $\beta$  receptors ALK4, ALK5, and ALK7 and activates the SMAD signaling pathway (Ho et al.). GDF-11 regulates the development of the olfactory system, retina, and pancreas, as well as anterior/posterior patterning of the axial skeleton (Lee & Lee). GDF-11 is an endocrine factor expressed in skeletal muscle, brain, and dental pulp (Kondás et al.). Studies in mice showed that GDF-11 regulates muscle and cardiac aging, and stimulates neurogenesis by remodeling blood vessels (Katsimpardi et al.; Loffredo et al.; Sinha et al.). This product is animal component-free.

## Product Information

**Alternative Names:** BMP-11, Bone morphogenetic protein 11, Growth/differentiation factor 11  
**Accession Number:** O95390  
**Amino Acid Sequence:** NLGLDCDEHS SESRCCRYPL TVDFEAFGWD WIIAPKRYKA NYCSGQCCEYM FMQKYPTHHL VQQANPRGSA GPCCTPTKMS PINMLYFNDK QQIYGGKIPG MVVDRCGCS  
**Predicted Molecular Mass:** 12.5 kDa monomer; 24.9 kDa dimer  
**Species:** Human  
**Cross Reactivity:** Mouse, Rat  
**Formulation:** Lyophilized from a sterile-filtered solution containing 0.1% trifluoroacetic acid.  
**Source:** E. coli

## Specifications


**Activity:** The specific activity is  $\geq 1.0 \times 10^4$  units/mg ( $EC_{50} \leq 100$  ng/mL) as determined by an alkaline phosphatase activity assay using ATDC-5 cells.  
**Purity:**  $\geq 95\%$   
**Endotoxin Level:** Measured by kinetic Limulus amoebocyte lysate (LAL) analysis and is  $\leq 1$  EU/µg protein.

## Preparation and Storage

**Storage:** Store at -20°C to -80°C.  
**Stability:** Stable as supplied for 12 months from date of receipt.  
**Preparation:** Centrifuge vial before opening. Reconstitute the product in 10 mM hydrochloric acid to at least 0.1 mg/mL by pipetting the solution down the sides of the vial. Do not vortex.

OPTIONAL: After reconstitution, if product will not be used immediately, dilute with concentrated bovine serum albumin (BSA) to a final BSA concentration of 0.1%. The effect of storage of stock solution on product performance should be tested for each application. As a general guide, do not store at 2 - 8°C for more than 1 month or at -80°C for more than 3 months. Avoid repeated freeze-thaw cycles.

## Data



(A) The biological activity of Human Recombinant GDF-11, ACF was tested by its ability to induce alkaline phosphatase activity in ATDC-5 cells. Alkaline phosphatase activity was measured using a fluorometric assay method. The EC<sub>50</sub> is defined as the effective concentration of the growth factor at which alkaline phosphatase activity is at 50% of maximum. The EC<sub>50</sub> in the example above is 3.54 ng/mL.

(B) 1 µg of Human Recombinant GDF-11, ACF was resolved with SDS-PAGE under reducing (+) and non-reducing (-) conditions and visualized by Coomassie Blue staining. Human Recombinant GDF-11, ACF has a predicted molecular mass of 24.9 kDa (12.5 kDa per monomer).

## Related Products

For a complete list of cytokines, as well as related products available from STEMCELL Technologies, visit [www.stemcell.com/cytokines](http://www.stemcell.com/cytokines) or contact us at [techsupport@stemcell.com](mailto:techsupport@stemcell.com).

## References

- Ho DM et al. (2010) The role and regulation of GDF11 in Smad2 activation during tailbud formation in the *Xenopus* embryo. *Mech Dev* 127(9–12): 485–95.
- Katsimpardi L et al. (2014) Vascular and neurogenic rejuvenation of the aging mouse brain by young systemic factors. *Science* 344(6184): 630–4.
- Kondás K et al. (2008) Both WFIKK1 and WFIKK2 have high affinity for growth and differentiation factors 8 and 11. *J Biol Chem* 283(35): 23677–84.
- Lee Y-S & Lee S-J. (2013) Regulation of GDF-11 and myostatin activity by GASP-1 and GASP-2. *Proc Natl Acad Sci USA* 110(39): E3713–22.
- Loffredo FS et al. (2013) Growth differentiation factor 11 is a circulating factor that reverses age-related cardiac hypertrophy. *Cell* 153(4): 828–39.
- Sinha M et al. (2014) Restoring systemic GDF11 levels reverses age-related dysfunction in mouse skeletal muscle. *Science* 344(6184): 649–52.

STEMCELL TECHNOLOGIES INC.'S QUALITY MANAGEMENT SYSTEM IS CERTIFIED TO ISO 13485. PRODUCTS ARE FOR RESEARCH USE ONLY AND NOT INTENDED FOR HUMAN OR ANIMAL DIAGNOSTIC OR THERAPEUTIC USES UNLESS OTHERWISE STATED.

Copyright © 2017 by STEMCELL Technologies Inc. All rights reserved including graphics and images. STEMCELL Technologies & Design, STEMCELL Shield Design, and Scientists Helping Scientists are trademarks of STEMCELL Technologies Canada Inc. All other trademarks are the property of their respective holders. While STEMCELL has made all reasonable efforts to ensure that the information provided by STEMCELL and its suppliers is correct, it makes no warranties or representations as to the accuracy or completeness of such information.